

A FEW COMMENTS ON CLASSICAL ELECTRODYNAMICS

K. Ghosh*

*Department of Engineering Science, Haldia Institute of Technology,
Hatiberia, Purba Midnapore - 721 657, India.*

I. ABSTRACT

In this article we discuss a few aspects of the space-time description of fields and particles. In section II we will show that a line is not just a collection of points and we will have to introduce one-dimensional line-intervals as fundamental geometric elements. Similar discussions are valid for area and volume-elements. In section III we will discuss the fundamental nature of fields. In section IV we will discuss non-equivalence of the Schwarzschild coordinates and the Kruskal-Szekeres coordinates. In App.B and App.C we make a comparative study of Quantum Field Theory and Quantum Mechanics and disagreements associated with probabilistic interpretation of these theories with space-time dimensional analysis. In App.D and App.E we discuss the geometry of Robertson-Walker model and electrostatic behavior of dielectrics respectively. In Supp.I we discuss the regularity of Spin-Spherical harmonics. In Supp.II we will give a causal description of interference and diffraction in terms of phase velocity of waves. In Supp.III we will discuss the regularity conditions required for the integral version of Gauss's divergence theorem and rederive Gauss's law in Electrostatics. We also show In Supp.IV we derive the complete equation for viscous compressible fluids and make a few comments regarding some contradictions associated with boundary conditions for fluid dynamics. In Supp.V We make with a few questions in We make a few comments on seeing by human brains in Supp.VI. We make a few discussions on radiation in isothermal enclosures and uncertainty principle in Sup.VII. In Supp.VIII We make a few discussions on Newton's laws for conservative force fields. In Supp.IX we derive an expression for scalar field entropy in the Schwarzschild black-hole background using the brick-wall model which is different from that of ' Hooft. We conclude this article with a few discussions on surface integral of vector fields in the last section "Miscellaneous".

* E-address: kaugho@rediffmail.com

II. GEOMETRIC CONTINUITY AND COORDINATIZATION

In this section we will review a few basic concepts of geometry and coordinatization. We will first give a precise definition of geometric continuity based on the concepts of homogeneity and extension. We will demonstrate that a line-element, a geometric continuum of one-dimension can not be a collection of points. Line-elements, area-elements and volume-elements are as fundamental as points. It follows naturally that an area-element can not be described as a collection of line-elements. This leads to a reformulation of the definition of dimension for a geometric continuum. We will consider the fact that real numbers form a dense set: for real numbers only the concept of "two numbers separated by an interval" or "two numbers are same" have meaning but "two numbers adjacent" is not defined. We will illustrate the relationship of these two aspects in the context of coordinatization. We will also state a corresponding version of the completeness of the real numbers. The definition of geometric continuity as given in this article raises the question whether time is a geometric continuum. We will briefly address this issue through the conceptual foundations of the General Theory of Relativity. We will also discuss in brief the the relationships of these aspects and the three spatial dimension with the the kinematical and dynamical properties of the Elementary particles and Electromagnetic fields. We will conclude this section with a few consequences.

In this section we will demonstrate that a line-element is not a collection of points. Line-elements, area-elements and volume-elements are as fundamental as points. We will consider the proposition similar to the fact that rational numbers form a dense set: for real numbers only the concept of "two numbers separated by an interval" or "two numbers are same" have meaning but "two numbers adjacent" is not defined. We will illustrate the relationship of these two aspects in the context of coordinatization.

The concept of geometric continuity should be based on two aspects: extension and homogeneity (more properly uniformity). An elementary particle, a tree if we neglect microscopic non-homogeneities, space and space-time (we will illustrate this example later) are examples of geometric continuum. In the most broad sense geometric continuity is distinguished according to dimension. An elementary particle, an electron, is a three dimensional geometrical continuum. If we consider a plane passing through the center of the electron (an equatorial cross-section) and shrink the transverse geometric extensions to zero we get a two-dimensional geometric continuum: a disk. If we consider a diameter of the disk and shrink the transverse extensions along the disk to zero we get a one-dimensional geometric continuity: a line. This definition of dimension is different from the conventional definition in terms of possible linear motion of point particles [32] but is proper as we will illustrate later. A point is a geometric object with zero extension. Thus whatever may be the amount of content (cardinality), a collection of points can not give a geometric continuum, a geometric object with finite homogeneous extension. We can illustrate this aspect in the context of the infinite straight line, a geometric continuum of one dimension. The one-dimensional homogeneous extension of a line-element on the straight line is characterized through the concept of length. Qualitatively, length characterizes finite homogeneous extension in one-dimension and also ordering of points with respect to a given point chosen as origin. Quantitatively, length gives a coordinatization scheme characterizing line-intervals with a given line-interval chosen as unity. We will illustrate this aspect later. Thus, whatever may be the cardinality, a collection of zero-length points can not give a finite-length one-dimensional line. Also a homogeneous collection of points give us a single point as the length of a point is zero. We can illustrate this aspect in the following way:

We consider a homogeneous linear array of marbles touching each other. If we now shrink the volume of each marble keeping them in contact (so that the array is always homogeneous) then in the limit that the volume of the marbles is zero we will get a single point.

These discussions lead us to conclude that a line-element is not just a collection of points but is a fundamental geometric object. Similar arguments lead us to conclude that area-elements (two-dimensional geometric continuums) and volume-elements (three-dimensional geometric continuum) are fundamental geometric objects and we can not obtain points without breaking the continuum geometry. To illustrate, let us consider a square, a geometric continuum of two dimension. We can not consider the square to be a collection of straight lines parallel to the base as each straight line has zero transverse extension and arguments similar to as above apply.

We will now discuss the concept of coordinatization which is an ordering process through length with respect to a point chosen as the origin. In the conventional scheme the infinite straight line is considered to be a collection of points and real numbers should have a one-to-one relationship with the points to have a proper ordering. In the context of the discussions in the preceding paragraph this scheme can not work as we can not arrange, according to length from a given point, a homogeneous collection of points each of which is of zero length. We have also illustrated earlier that a homogeneous collection of points give us a single point. A plane is a two dimensional object and any point on the plane can be characterized, as we will illustrate, through the choice of two intersecting lines on the plane and constructing a coordinate system

defined in terms of intervals with respect to the point of intersection.

We now consider the concept of coordinatization more precisely. As we have discussed earlier line intervals as fundamental geometrical entities to form lines which are geometric continuum of one dimension.

We define a dense set to be a set with an ordering such that there exist an element between any two distinct elements of the set. We will now prove that the real numbers form a dense set [33,34]. That is for real numbers only the concept of "two numbers separated by an interval" or "two numbers coincident" have meaning but "two numbers adjacent" is not defined. That is for any two real numbers we can always find a real numbers lying between the two.

Firstly, the two representations of 0.1, 0.0999..... (where all the decimal places starting from the 2-nd are 9) and 0.1 are same as any number added to 0.0999..... gives a number greater than 0.1 and any number subtracted from 0.1 gives a number less than 0.0999..... This two representations also gives us the same result if add/subtract a number with/from both the numbers.

An arbitrary real rational number may be expressed as $r = n.x_1x_2x_3.....x_p$, where n is an integer, $x_i = 0, \dots, 9$ for $i < p$, $x_j = 0$ for $j > p$, p may be arbitrarily large but finite and $x_p \neq 0$, i.e., the sequence of the decimal places is finite. Whereas an irrational number and a recurrent decimal is given by: $ir = n.x_1x_2x_3.....$ where all the decimal places are significant. We will consider $r, ir > 0$ in the following section.

We now prove the following proposition:

For real numbers only the concept of "two numbers separated by an interval" or "two numbers coincident" have meaning but "two numbers adjacent" is not defined. That is given any two real numbers we can always find a real numbers lying between the two.

If r is an integer (> 0) and $M = r.x_1x_2x_3.....x_p$, with all but $x_p (= 1)$ are zero and p may be arbitrarily large, then the number of decimal places having value zero before $x_p = 1$ can be increased arbitrarily to construct a number less than M and closer to r and it is not possible to define a number to be adjacent to r . Similar will be the case for $N = (r - 1).999.....x_i$, with all the decimal places up to x_i are 9 for i arbitrarily large, to construct a number greater than N and closer to r , i.e, the number of decimal places having value 9 can be increased arbitrarily and it is not possible to define a number to be adjacent to r . We can illustrate this issue with the representations discussed before the proposition. The numbers r and $(r - 1).999...$ where all the decimal places are 9 equivalent. Thus we can form the number $(r - 1).999.....x_i = 899..$ and applying arguments similar as above to the i -th decimal place we can show that it is not possible to define a number adjacent to r . This is also same as subtracting $M = 0.000.....x=1$ from r and applying arguments similar to those discussed at the beginning of this paragraph.

Similar as above will be the arguments with ($n \geq 0$) $r = n.x_1x_2x_3.....x_p$, $M = n.x_1x_2x_3.....x_j$ (x_i same for both for $1 \leq i \leq p$) with all x_i ($i > p$) but $x_j (= 1)$ ($j > i$) are zero for j arbitrarily large and for $N = n.x_1x_2x_3.....y_p999.....x_j$ with the p -th decimal place $y_p = x_p - 1$ and all x_j for j arbitrarily large are equal to 9.

For irrational numbers like $(ir)_1 = \sqrt{2}, \pi$ we can prove the proposition using the rational approximation as considered in the standard literature. We can also apply the above arguments for recurrent decimals. Following the constructions in the preceding paragraphs it is easy to show that between any two irrational number there exist a rational or irrational number.

For $r \leq 0$ the corresponding arguments to prove that "two numbers are adjacent" is not defined are very similar as in the preceding paragraphs.

These arguments together with the completeness of the Decimal number system (in the context of number systems) prove the above proposition.

We now consider a few aspects about irrational numbers defined in the conventional way. If we represent irrational numbers in terms of nested families of closed intervals [32] with rational boundaries then we will have to assign a corresponding non-zero length (as for Euclidean space length is same as the difference of the rational boundaries) to the null rational interval representing a single point on the real line corresponding to the irrational number and this is not well-defined. On the other hand we can assume that between two rationals there exist a set of irrational numbers with forming a line-interval with continuum cardinality [6]. (This has some consequence for first countability of the real line as we will discuss at the end of this section) We now consider the null rational interval $[0, r]$. Whatever r can be we can multiply this interval with a factor of the form $(10)^M$ (r can be a surreal number and M can be a superreal number [35]). The resultant interval should contain an infinite number of rationals [33] although all the numbers in the interval $[0, r]$ (apart from the two boundaries) are irrational and multiplying an irrational number by $(10)^M$ can not give us a rational number. Thus we take the point of view of [36], i.e, the rational approximations of an irrational number form a dense set about the irrational number and the rational approximation give us the irrational number in the limit when the number of significant decimal places are infinite, i.e, when the number of significant decimal places never terminate. For a recurrent decimal and an irrational to be adjacent the

irrational should differ from the rational at some finite decimal place and arguments similar as discussed earlier demonstrate that it is not possible to define an irrational and a recurrent decimal to be adjacent.

These discussions also lead us to conclude that it is not possible to define a successor for real numbers. If we do not accept the point of view of [34] to break the fact that real numbers form a dense set then this aspect of real numbers is in accordance with geometric continuity as defined at the beginning of this article. That is, as far as coordinatization is concerned, the real numbers characterize line-intervals, which can be arbitrary, with respect to a point chosen to represent the value zero, the origin. This scheme of coordinatization differs from the conventional way where we consider the infinite straight line to be a collection of points with continuum cardinality and try to have a one-to-one relationship between the points and the real numbers. In this context we can reformulate the completeness of the real numbers in the following way:

If we chose any interval on the infinite straight line (the Real line in the sense that the intervals on the line is characterized by real numbers with a point on the line chosen to represent the number zero) the length can be expressed either in terms of a rational number or an irrational number. Similar will be the situations with the higher dimensional volume elements.

We note as the real line cannot be defined as a collection of points we have to introduce the concept of collection of one-dimensional line elements as a fundamental mathematical entity which can be characterized into four classes:

- (i) one-dimensional line elements without boundaries
- (ii) one-dimensional line elements with one boundaries
- (iii) one-dimensional line elements with two boundaries
- (iv) one-dimensional line elements with the two boundaries identified (e.g, a circle)

We now demonstrate the consistencies of the discussions regarding geometric continuity at the beginning of this article in the following way. We can define the radius of a circle as the perimeter over 2π . As is proved earlier the values of the radius can only be defined through intervals and the concept of two concentric circles with adjacent values of radii is not defined. This feature is consistent with the fact that we can not obtain a two-dimensional disk from a collection of one-dimensional circles. We have to introduce the two dimensional circular strips together with the point origin as fundamental mathematical objects to construct a two-dimensional disk in the above way. Similar arguments for two dimensional spheres (where the radius is now defined as the positive square root of the area of the two-dimensional sphere over 4π) illustrate that we can not have a three dimensional volume element from a collection of two dimensional spheres and we have to consider three dimensional volume elements as fundamental mathematical entities together with the point origin. These discussions can be extended to higher dimensions.

A natural consequence of the above discussions is the fact that even if a physical variable is a geometric continuum its values are defined through intervals with respect to a reference value and form a dense set. Existence of quanta are restrictions on the values of these intervals leading to isolated spectrum.

The finite volume of the elementary particles and their displacements are in accordance with the geometric continuum structure of space. Within the context of classical mechanics time is realized through the changes in the configuration of a system. To illustrate we can consider the coordinates of a moving particle to characterize time. In the conventional definition of a line as collection of points we can define time to be a geometric continuum, as a collection of points, by a one-to-one correspondence between the possible coordinates of the particle on the line and the instants of time. However as we have discussed in this article a line is not a collection of points. Line-intervals are fundamental geometric objects. Only changes of position of the particle by a non-zero line-element is meaningful. The coordinates do not form a geometric continuum but form a dense set characterizing line-intervals with respect to a chosen point as the origin. Thus if we define the instants of time through a one-to-one correspondence with the possible coordinates of the moving particle we will have a dense set structure for time. However the General Theory of Relativity is a theory governing the dynamics space-time intervals themselves. The kinematical equivalence of space and time intervals indicates that time should be a geometric continuum in the same way that space is. The space-time coordinates of events form a dense set.

Unlike the motion of points, line-elements and area-elements the motion of three-dimensional objects form a geometric continuum of the same dimension as that of the objects themselves. Thus the three dimensional spatial geometry of the Universe is realized through the finite three dimensional volume of the fundamental particles and the finite three dimensional volumes of the fundamental particles lead to three spatial dimensions for the Universe. Dynamically, there are two possible motions on a plane, translations and rotations. Dynamically infinitesimal rotations can be uniquely characterized through their magnitude and the direction of the normal to the plane of rotation chosen through proper convention. In three dimensions the normal to the plane of infinitesimal rotation is uniquely characterized up to orientation, determined by proper convention, and is contained within the three dimensional spatial geometry. Higher dimensions greater than three break the uniqueness of the normal to the plane of rotation. Thus three dimensional spatial

geometry is self-complete to describe the dynamics of matter particles uniquely. The quantum mechanical description of spin and polarization of the elementary particles is also significant in this respect. Also the concept of orientation with a proper convention is an essential aspect to formulate the laws of Classical Electrodynamics consistent with the energy conservation law [24]. The fact that in general, apart from the case when the normal to the plane of the loop is perpendicular to the magnetic field, a time-varying magnetic field always induces an electric current in a closed loop obeying Faraday's law (consistent with the energy conservation law) indicates that spacial geometry is of three dimensions as far as Classical Electrodynamics is concerned.

To summarize, in this section we have established that a line-interval is not a collection of points and we will have to consider line-intervals as fundamental geometric objects. Similar discussions are valid for higher dimensional geometric continuum. We have used the fact that real numbers form a dense set: "For real numbers only the concept of "two numbers separated by an interval" or "two numbers coincident" have meaning but "two numbers adjacent" is not defined" and discussed the relation of these two facts in the context of how to define coordinatization. We have also discussed in brief the the relationships of these aspects and the three spatial dimension with the the kinematical and dynamical properties of the Elementary particles and Electromagnetic fields.

We conclude this section with a few discussions:

Firstly, we can not express the Cartesian plane R^2 in terms of the direct product $R^1 \times R^1$. R^1 is not just a collection of points. Hence it is not possible to express R^2 as the union of the corresponding number of R^1 s. Also we have discussed earlier that we can not express a two-dimensional geometric continuum as a homogeneous collection of one-dimensional geometric continuum. Since R^2 can not be expressed as a collection of curves, the tangent vector space at any point on R^2 can not be identified with R^2 . The above discussions are also valid in higher dimensions. This aspect also illustrate that, we can not define higher dimensions greater than one in terms of possible motion of a point particle as neither a two-dimensional area-element nor a three-dimensional volume-element is a collection of one-dimensional line-elements. These discussions are significant for Differential Geometry. We now illustrate in brief the significance of these discussions in the context of classical statistical mechanics of a system of particles. We define the configuration-space to be the collection of the spatial coordinates of the particle. As we have illustrated earlier the configuration-space forms a dense set (a dense vector space when the dimension is greater than one). The momentum of a particle moving along a line is defined through associating possible values of the momentum with the possible positions of the particle in a given time starting from a fixed point. For a single particle moving in one-dimension the corresponding momentum at any point is not a geometric continuum but form a dense set as changes in the position of the particle through arbitrary but non-zero intervals are only meaningful. This discussions are similar to those made for time. Also the discussions at the beginning of this paragraph indicates that in higher dimensions the momentum of the particle at any point, tangent vector to the possible trajectories of the particle at that point, do not form a geometric continuum but is a dense vector space. Thus the dynamical states of the particle is characterized by the coordinate dense set (the configuration-space) and a dense collection of the momentum space dense set. The fact that the coordinates and the momentums form a dense (vector) space indicates that physically we can only approximate the corresponding phase-space to a geometric continuum. The above discussions can be extended when the number of particles is greater than one. We will later illustrate the significance of these discussions in the context of the corresponding path-integral formulation of statistical mechanics. In passing we note the following: if we express a geometric continuum as a collection of points the cardinality and hence the amount of accessible microstates remains to the same irrespective of dimension. It is through coarse graining that one obtains different expressions for entropy depending on dimension.

Secondly, let us consider two sets each containing a single object: a closed line-element. The line-elements are intersecting but non-coincident everywhere. The intersection of these two sets is a set containing points which are fundamentally different from line elements. Also as a line-element is not just a collection of points a set-theoretic union to form a closed line-element out of a set of half-open line-elements and a set of points is not well-defined as the above mentioned sets contains elements that are geometrically and thereby intrinsically different. Similar discussions remain valid in two dimensions when two area elements intersect along a line-element and in three dimensions when two closed volume-element intersect along a common boundary. These problems may be solved by defining a universal set containing all possible geometric-elements including points. The union or intersection of any two elements from this universal set give us elements which belong to this universal set.

Lastly we discuss a consequence of the discussions on irrational numbers as discussed earlier.

To illustrate we make some comments regarding the first and second countability of R^1 discussed in Appendix:A of Wald [3]. A topological space X is first-countable if for every point p belonging to X there is a countable collection of open sets such that every open neighbourhood of p contains at least one member

of the collection. Whereas X is second countable if there is a countable collection of open sets such that every open set can be expressed as an union of open sets from this family. For R^1 , the open balls with rational radii centered on points with rational coordinates form such a countable collection open sets. When defined as in [6] there are infinite number of irrational numbers between two rational numbers. As far as first-countability is concerned, an open set V centered on a rational number with deleted perimeter on one of these neighbouring irrational number cannot contain open balls with rational radii centered on points with rational coordinates. Similar discussions can also be extended to R^n .

III. FUNDAMENTAL NATURE OF FIELDS

Gravitational interaction which is universally attractive is described by the General Theory of Relativity, a theory of fields realized through the existence of space-time and some of its geometric properties, e.g., the curvature. The principle of equivalence leading to the fact that the guinea and the feather fall the same way in vacuum together with the fact that gravity violates the elementary quantum mechanical principles (as will be evident from this article) indicates that fields are as fundamental as particles. Fields and particles with their common and contradictory kinematical and dynamical features are the two fundamental constituents of nature. The attraction of opposite charges can only be explained through accepting the electric field as a fundamental entity of nature and not through only interaction mediating particle interaction (be the interaction mediating particles interacting with the sources or among themselves) although the repulsion between like charges may be explained in the later way with photons as the interaction mediator. Quantum mechanics is an incomplete approach to explain the Solar system microscopic physics purely in terms of wave-like properties whereas Quantum Field Theory is an incomplete approach to explain the Solar system microscopic physics purely in terms of particle-like properties. For the vacuum polarization explanation of the potential [18] in the Bhaba scattering process it is not obvious how, in terms of photon exchanges, the loops in vacuum will screen the charges of e^- , e^+ . Also an electron-positron loop with two external photon lines can not explain charge screening in any process as the particle-antiparticle pairs are created and annihilated after the interaction mediating photon had been created at one real particle and before the interaction mediating photon has interacted with the other real particle. These two theories are also contradictory to gravity (a theory of space-time) in many aspects as will be manifested in this article.

During the last few decades a lot of efforts had been devoted to unify the general theory of relativity (describing the gravitational interaction) with quantum mechanics (describing the microscopic interactions of the elementary particles). Yet the conventional theory of quantum mechanics, based on unitarity and symmetries, is contradictory with general relativity in many respects, *e.g.*, formally infinite zero point energy associated with canonical quantization scheme, ultraviolet divergent energy density associated with vacuum fluctuations [1] for collapsing physical systems, unitarity violation for black hole evolution.

To unify these two descriptions of nature we can proceed along two directions [2] :

For the Euclidean school space-time geometry is an abstract concept which exist irrespective of matter fields. For the non-Euclidean school space-time does not exist independent of matter fields. Space-time is form of existence of matter and can not be conceived without matter. This feature is even more transparent from the facts, among others, that the universe is compact and there is no well-defined stress-tensor for gravity which intertwinds matter fields and space-time geometry [3].

In the context of the general theory of relativity the conflict between the two schools arise in the following way:

As soon as one derives the geodesic deviation equation from the principle of equivalence one “can” forget the source, ascribing the relative acceleration of the nearby geodesics to the space-time manifold. This, in contradiction to the philosophy of general relativity, may lead to think that the space-time manifold is more fundamental leading to the concept of quantum gravity irrespective of existence of the corresponding sources that will produce the fluctuating geometries (no source indicates no space-time geometry and observationally, quantum fluctuations in matter fields are negligible to produce significant alternations of the space-time geometry).

Quantum gravity also led to many space-time geometries which are physically non-existent. One such example is the extreme Reissner-Nordstrom black hole which cannot be obtained through any realistic gravitational collapse []. It also has vanishing Hawking temperature. This may be interpreted through the fact that Hawking radiation through pair production near the black hole event horizon is not possible as the metric do not change its signature across the horizon.

We should keep in mind that the Einsteins equations in the regime of its validity determine the space-time geometry: the geometry of space-time as a whole is determined by corresponding matter fields described either in terms of some classical models or by a proper quantum theory. This gives a particular cosmology

(the closed or mathematically more properly compact universe picture) and if we view cosmology as a whole there is really no test body.

We will now consider some aspects of the black hole space-time geometry. In the process of gravitational collapse an event horizon, the black hole event horizon, is formed breaking the global CP invariance and giving rise to the Kerr-Newman families of black holes (the no hair theorems). The black hole event horizon may be defined as the causal boundary of the set of complete time-like geodesics which originates at the past time-like infinity and terminate at the future time-like infinity as classically nothing can come out off the horizon. The black hole space-time is usually described either in terms of the Schwarzschild coordinate system or in terms of the Kruskal-Szekers coordinate system. In the Schwarzschild coordinate system the black hole event horizon is a two dimensional fixed point set of the time-like Killing vector field across which some of the metric components change sign. In the Kruskal-Szekers coordinate system the event horizon is a two dimensional null surface across which the square of some of the coordinates change sign. We will now consider the non-equivalence of the two coordinate systems in detail.

IV. NON-EQUIVALENCE OF THE SCHWARZSCHILD AND THE KRUSKAL-SZEKERS COORDINATE SYSTEM

The Schwarzschild space-time is a Lorentz signature, static spherically symmetric solution of the Einstein equations when the Ricci tensor vanishes. This solution describes the exterior geometry of a static spherically symmetric star and has been used to verify the predictions of general relativity for the Solar system.

A space-time is said to be static if there exists a space-like hypersurface which is orthogonal to the orbits of the time-like Killing vector field. A space-time is said to be spherically symmetric if the space-like hypersurfaces contains $SO(3)$ as a subgroup of the group of isometries. The orbit spheres of $SO(3)$ are isometric to the unit two sphere. These features together with the condition of the asymptotic Newtonian limit give the well-known Schwarzschild solution in the spherical polar coordinates[3]:

$$ds^2 = -(1 - 2M/r)dt^2 + (1 - 2M/r)^{-1}dr^2 + r^2[d\theta^2 + \sin^2\theta d\phi^2] \quad (1)$$

According to the Birkhoff's theorem [19] all spherically symmetric solutions with $R_{ab} = 0$ are static and the Schwarzschild space-time is the unique static spherically symmetric solution, up to diffeomorphisms, of the Einstein equations with $R_{ab} = 0$.

The norm of the time-like Killing vector field and $(\nabla r)^a$ in the orthonormal coordinates vanishes and some of the metric components are not well-behaved at $r = 2M$ in the Schwarzschild coordinates. The proper acceleration of the constant r observers can be obtained from the geodesic equations in the Schwarzschild coordinates. This acceleration, $a = (1 - 2M/r)^{-1/2}M/r^2$, is divergent at the horizon ($r = 2M$).

The ill-behavedness of the Schwarzschild coordinates is not a coordinate singularity like that of the spherical polar coordinate system where the azimuthal angular coordinate ϕ become ambiguous at the poles. All the ill-behavedness of the Schwarzschild coordinates at the horizon originate from that of the space-time metric. The curvature scalars calculated from the metric are well-behaved at the horizon unlike $r = 0$ where the curvature scalars diverge. For ordinary stars this metric singularity at $r = 2M$ is irrelevant as it is inside the star and the Schwarzschild solution is not valid in the matter filled interiors. However it is well-known that sufficiently massive stars can undergo gravitational collapse to form black holes and the metric singularity at the horizon is important. Several coordinate systems had been introduced to remove the metric singularity and to extend the Schwarzschild space-time where the Schwarzschild coordinate system is referred to covering a proper submanifold of the extended space-time. The metric in these extended coordinate systems are well-defined every where apart from the space-time singularity. The most well-known extension is the Kruskal-Szekers coordinate system. In this article we perform a comparative study of these two coordinate systems and show that they are not diffeomorphically equivalent.

In this section we will follow the abstract index convention of Wald [3] and extend its significance in Appendix:A.

According to the theory of relativity if $\phi : M \rightarrow M$ is diffeomorphism then (M, g_{ab}) and $(M, \phi^* g_{ab})$ represent the same physical space-time. Let a coordinate system x^μ cover a neighborhood U of a point p and a coordinate system y^ν cover a neighborhood V of the point $\phi(p)$. Now we may use ϕ to define a new coordinate system x'^μ in a neighborhood $O = \phi^{-1}[V]$ by setting $x'^\mu = y^\mu[\phi(q)]$ for q belonging to O . We may then take the point of view as ϕ leaving p and all tensors at p unchanged but inducing the coordinate transformation $x^\mu \rightarrow x'^\mu$. For ϕ to be a diffeomorphism $\frac{\partial x'^\mu}{\partial x^\nu}$ should be non-singular [3,15]. According to this point of view two coordinate system covering a space-time can be taken to be equivalent if the corresponding

transformation coefficients are not singular in their common domain of definition otherwise an arbitrary smooth function defined in one coordinate system may not remain smooth in the other coordinate system.

To extend the Schwarzschild coordinate system one considers the two dimensional $r - t$ part:

$$ds^2 = -(1 - 2M/r)dt^2 + (1 - 2M/r)^{-1}dr^2 \quad (2)$$

The Regge-Wheeler coordinate system is defined through the null-geodesics and is given by:

$$r_* = r + 2M \ln(r/2M - 1) \quad (3)$$

in this coordinate $r \rightarrow 2M$ corresponds to $r_* \rightarrow -\infty$. The null coordinates are defined as:

$$u = t - r_*, \quad v = t + r_* \quad (4)$$

A regular metric is obtained through the following transformation,

$$U = -e^{-u/4M}, \quad V = e^{v/4M} \quad (5)$$

The metric in these coordinates becomes:

$$ds^2 = -\frac{32M^3 e^{-r/2M}}{r} dU dV \quad (6)$$

As there is no longer a coordinate singularity at $r = 2M$ (i.e at $U = 0$ or $V = 0$) one extends the Schwarzschild solution by allowing U, V to take all possible values. However the transformation coefficients $dU/dr = -d[(r/2M - 1)^{1/2} e^{-\frac{(t-r)}{4M}}]/dr$ and $dV/dr = d[(r/2M - 1)^{1/2} e^{\frac{(t+r)}{4M}}]/dr$ are singular at $r = 2M$ and the extension is not diffeomorphically equivalent. Consequently as discussed at the beginning of this section the Schwarzschild coordinate system and the (U, V) coordinate system do not represent physically the same space-time manifold. Consequently, according to Birkoff's theorem, the space-time represented by the (U, V, θ, ϕ) coordinate system is not a solution of the Einstein equations for a spherically symmetric black hole.

Similar discussions are valid for the Kruskal-Szekers coordinate transformations which are obtained through the following transformations:

$$T = (U + V)/2, \quad X = (V - U)/2 \quad (7)$$

and the metric becomes,

$$ds^2 = \frac{32M^3 e^{-r/2M}}{r} (-dT^2 + dX^2) + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \quad (8)$$

The relation between the (T, X) and the (t, r) coordinates are well known and in the physical regions of interests are given by [4],

$$X = (r/2M - 1)^{1/2} e^{r/4M} \cosh(t/4M) \quad (9)$$

$$T = (r/2M - 1)^{1/2} e^{r/4M} \sinh(t/4M) \quad (10)$$

valid for $r > 2M$, and

$$T = (1 - r/2M)^{1/2} e^{r/4M} \cosh(t/4M) \quad (11)$$

$$X = (1 - r/2M)^{1/2} e^{r/4M} \sinh(t/4M) \quad (12)$$

valid for $r < 2M$.

Again the transformation coefficients are not defined on the horizon and the Kruskal-Szekers coordinates do not give a proper diffeomorphic extension of the Schwarzschild coordinate system. Hence the Kruskal-Szekeres coordinates is not a solution of the Einsteins equations for a spherically symmetric black hole

The Kruskal-Szekers coordinate system had been introduced to eliminate a particular singular function (the metric components) in the Schwarzschild coordinate system through a singular coordinate transformation. This does not ensure that all singular tensors can be made regular in the new coordinate system and also tensors which are regular in the (t, r) coordinates can become singular in the (T, R) coordinates. To illustrate these features we consider the implicit relations between the two coordinate systems [1]:

$$(r/2M - 1)e^{r/2M} = X^2 - T^2 \quad (13)$$

$$\frac{t}{2M} = \ln\left(\frac{T+X}{X-T}\right) \quad (14)$$

The horizon in this coordinates are defined as $X = \pm T$.

Firstly the proper acceleration of the curves in Kruskal-Szecker's coordinate system which correspond to the constant r observers in the Schwarzschild coordinate system is given by $a = (X^2 - T^2)^{-1/2} [e^{r/2M} M/r^2]$. This is also divergent on the horizon.

Secondly we consider the vector $(\frac{dR}{ds})^a$, R'^a , the proper rate of change of the curvature scalar R obtained from $(dR)^a$ and the proper distance ds [i.e, the vector $(\frac{dR}{ds})(\frac{\partial}{\partial r})$, Appendix:A]. The norm of this vector in the Schwarzschild coordinate system is $(dR/dr)^2$ and is finite on the horizon. Whereas the corresponding quantity in the (T, X) coordinates can be obtained from the following relations [apart from normalizing factors: $(\frac{\partial X}{\partial s}), (\frac{\partial T}{\partial s}) = [\frac{re^{r/2M}}{32M^3}]^{1/2}$:

$$\frac{dR}{dX} = \frac{\partial R}{\partial r} \frac{\partial r}{\partial X}, \quad \frac{dR}{dT} = \frac{\partial R}{\partial r} \frac{\partial r}{\partial T} \quad (15)$$

and from equ.(13),

$$\frac{\partial r}{\partial X} = \frac{8M^2 X e^{-r/2M}}{r}, \quad \frac{\partial r}{\partial T} = -\frac{8M^2 T e^{-r/2M}}{r} \quad (16)$$

and we have $|R'^a_{KS}|^2 = \frac{64M^4 e^{-r/M}}{r^2} (\frac{\partial R}{\partial r})^2 [X^2 - T^2] = 0$ on the horizon although the r -dependent multiplying factor in front of the Kruskal-Szecker's metric is finite at $r = 2M$.

The unit space-like normal vector to the $r = \text{constant}$ surfaces, which can be defined apart from $r = 0$, $k^a = (\frac{dr}{ds})^a$ has unit norm ($k^a k_a = 1$) on $r = 2M$ although $k^a \rightarrow 0$ as $r \rightarrow 2M$ which for an outside observer ($r > 2M$) may be interpreted as nothing can propagate radially outward at $r = 2M$.

For two metric spaces the definitions of continuity is as follows [16]:

Let (S, d_S) and (T, d_T) be metric spaces and let $f : S \rightarrow T$ be a function from S to T . The function f is said to be continuous at a point p in S if for every infinitesimal $\epsilon > 0$ there is an infinitesimal $\delta > 0$ such that

$$d_T[f(x), f(p)] < \epsilon, \quad \text{whenever } d_S[x, p] < \delta. \quad (17)$$

If f is continuous at every point of S then f is continuous on S .

The definition is in accordance with the intuitive idea that points close to p are mapped by f into points closed to $f(p)$. From equn.(13),(14) we have,

$$|dt|_{Sch} = \frac{X}{(X^2 - T^2)^{1/2}} |dT|_{KS}, \quad |dt|_{Sch} = -\frac{T}{(X^2 - T^2)^{1/2}} |dX|_{KS} \quad (18)$$

and,

$$|dr|_{Sch} = \frac{X}{(X^2 - T^2)^{1/2}} |dX|_{KS}, \quad |dr|_{Sch} = -\frac{T}{(X^2 - T^2)^{1/2}} |dT|_{KS} \quad (19)$$

where $|\cdot|$ denotes the norm in the respective coordinate systems and we find that the coordinate transformation, $(t, r) \rightarrow (T, X)$ is not continuous on the horizon as the multiplicative factors diverge on the horizon ($X = \pm T$). Consequently the coordinate transformation $(t, r) \rightarrow (T, X)$ is not a homeomorphism and the two coordinate systems do not topologically represent the same space-time manifolds [3,17]. Hence we show that the Kruskal-Szekers coordinate system is not a proper extension of the Schwarzschild coordinate system and it is not a solution of the Einsteins equation for spherically symmetric black hole. We conclude this discussion with the following note:

For any coordinate system we have,

$$g'_{\mu\nu} = \frac{\partial x^\rho}{\partial x'^\mu} \frac{\partial x^\sigma}{\partial x'^\nu} (g_{Sch.})_{\rho\sigma} \quad (20)$$

Consequently it is not possible to find a coordinate system with a regular $g'_{\mu\nu}$ without absorbing the singularities of $(g_{Sch.})_{\rho\sigma}$ at $r = 2M$ into the transformation coefficients $\frac{\partial x^\rho}{\partial x'^\mu}$ at $r = 2M$ i.e, without breaking the diffeomorphic equivalence of the two coordinate systems. Thus, as also discussed in the preceding sections, the Kruskal-Szekeres coordinate system with a regular metric at the horizon can not be diffeomorphically equivalent to the Schwarzschild coordinate system and thus do not represent a static asymptotically flat solution of the Einsteins equations representing a black hole formed out of the gravitational collapse of an uncharged spherically symmetric asymptotically flat star. [see also Appendix:E].

It is not obvious how to describe the space-time evolution of the complete gravitational collapse of matter fields as a whole in terms of time-like curves as, for a Schwarzschild observer, the time-like curves suffer a discontinuity across the horizon and become space-like inside the black hole event horizon. It is wellknown that expressed in terms of the Schwarzschild coordinates the black hole event horizon has profound impact on the quantum description of matter fields and black hole evaporation through Hawking radiation makes the space-time dynamic. Also Hamiltonian evolution of matter fields break down on the fixed point sets of the time-like Killing vector field [14]. The canonically conjugate momentums are not well-defined on the horizon as will be evident from the lagrangians of the matter fields.

V. DISCUSSION

In a gravitational collapse once the collapsing body crosses the horizon it collapses to form the space-time singularity breaking the description of space-time in terms of continuous manifolds and the local symmetries. We can only characterize the presence of the space-time singularity in a diffeomorphism invariant way, in terms of the curvature invariants along the space-time curves which cross the event horizon and necessarily terminate along the space-time singularity. The formation of black hole event horizon can be characterized through the formation of trapped surfaces. The gravitational collapse and the cosmological evolution are the only two processes in nature through which a three dimensional physical system collapses to zero dimension (forming the space-time singularity). Here through zero dimension we mean a point or a collection of points. We will illustrate this aspect in Appendix:B.

Einsteins equations break down at the space-time singularity. This is something similar to electrodynamics. We can determine the electric field for a point charge using the Maxwells equations. But the field strength diverges and classical electrodynamics break down at the point charge (the corresponding quantum theory QED is not a resolution to this problem. It has its troubles associated with the point-like interaction terms. However experimental observations confirm that all the elementary particles are of finite volume). The formation of space-time singularity is associated with finite volume to zero dimension transition for the corresponding collapsing body and the richest structure that we can attribute to zero dimension is that of an analogue of (compact) three dimensional generalization of the Cantor set [5,6] provided we generalize the description of the collapsing matter field through a proper quantum theory [a generalization of the Pauli exclusion principle].

There are two ways that one can reach zero dimension from finite volume breaking the continuous topology of space-time manifold. One is through the point contraction mapping which requires an infinite number of iterations which, together with the discussions in Appendix:B, is in accordance with the fact that time is

a continuous parameter. The other one is through the formation of an analogue of the Cantor set (or any other discrete manifold with different cardinality) in which case the underlying physical process to achieve zero dimension may be discontinuous. A discrete manifold may not always form a normed vector space, e.g., the set of points $(n + x)$ on the real line, where n is an integer and x is a fractional number, can not form a normed vector space as the difference between two points do not belong to the set. Also it is not physically obvious to talk of causal structure, defined through propagation of signals, in a discrete manifold unless the manifold is space-like and frozen in time (which is defined through physical processes). As discussed earlier according to the General Theory of Relativity charges associated with space-time transformation symmetries are global properties of a continuous space-time manifold as a whole whereas a manifold without continuous topology can only have space-time independent charge.

To describe cosmological evolution and black hole evolution we will have to generalize and geometrize conventional quantum mechanics in a suitable way. In these respects the principal aspects to be critically studied, as will be discussed later in this article, towards unifying the general theory of relativity and conventional quantum mechanics is the concept of diffeomorphism invariance associated with the general theory of relativity and unitarity associated with the conventional quantum mechanics.

The facts that the continuous topology of space-time break down at the space-time singularity [Appendix:B] (indicating no well-defined observables associated with spatial transformations for the cosmologically evolving or collapsing matter fields in the near zero dimension region) and that nature chooses a particular cosmology lead us to conclude that diffeomorphism invariance (which for large scale structure of space-time is equivalent to invariance under coordinate transformations) is not of so important (as it is for solar system microscopic physics) concern for the corresponding physical laws. Rather the fact Schwarzschild coordinates and the Kruscal- Szekers coordinates are not diffeomorphically equivalent indicates that an appropriate choice of a suitable coordinate system is most important. However we can express the generalized quantum theory in covariant form. This will help us to compare the generalized quantum theory with solar system quantum physics where the physical laws are invariant under inertial transformations and are formulated in a covariant way under the corresponding coordinate transformations.

To generalize the conventional quantum mechanics we should take into account the following important aspects:

(1) Special relativity made the concept of size for ordinary objects a relative one. The strong curvature effects near the space-time singularity will spoil the concept of dimension for the elementary particles forming the space-time singularity.

(2) Quantum mechanics is the mechanics of quantum states which do not exist independent of their realizations at least in principle, i.e., through interactions with other quantum states. For solar system microscopic physics the fact that the elementary particles and bound states (e.g., atoms) formed by them are of finite volume has to be considered in the corresponding quantum state description as long as the continuous structure of the space-time manifold holds.

(3) Every measurement process through state reduction is a non-unitary operation on the space of quantum states [7]. Near the space-time singularity the strong curvature will destroy the description of matter fields in terms of a unitary quantum theory.

As far as the cosmological evolution is concerned (zero dimension to finite volume and finite volume to zero dimension transitions) no observer physics is the exact description of the evolution of the universe in the near zero volume region. A proper generalization of quantum mechanics may be non-unitary in the sense that the evolutions of the possible quantum states (if the space-time description of matter is given by a particular family of space-time curves representing possible particle trajectories [5]) representing the collapsing physical systems may be non-unitary.

However, it is obvious that the collapsing physical system (which is a bound system through gravitational interaction) collapses to zero dimension violating the conventional quantum mechanics based on the uncertainty principle (e.g., the electrons in an atom obeying the quantum mechanical principles do not collapse on the positively charged nucleus) and follow the deterministic laws of general relativity.

In the context of the above discussions an important contradiction between the general theory of relativity (describing the gravitational interaction) with conventional quantum mechanics (describing the microscopic interactions of the elementary particles) is the following:

Positivity of the energy momentum stress-tensor together with the general theory of relativity leads to gravitational collapses [8] and space-time singularities [9] where a three dimensional physical system collapses to zero dimension (breaking the continuous space-time topology) whereas positivity of the energy-momentum tensor together with the canonical commutation relations lead to the Pauli exclusion principle (unless one introduces additional structures about the space-time singularity).

We, living beings, are characterized by the fact that we can control some terrestrial processes. But we can neither change the physical laws nor the cosmological evolution. Many descriptions we had made are

either approximations (unitary quantum mechanics, quantum statistical mechanics) or idealizations (point particles for microscopic physics). The discussions in the preceding paragraph together with the facts that black holes contain no scalar hair [11], that there is no physical explanation of only recombination for the virtual particles (which “interpret” real effects as in the Casimir effect) to form loops in quantum field theory [Physically, in vacuum, even in Feynmann’s summing over path scheme it is not obvious why particle pairs produced at one space-time point will only recombine at another space-time point. The otherwise should give abundance of particles and antiparticles. We will discuss this issue in some details in Appendix:C.] and that the universality of the minimum uncertainty relations are lost in the gravitational collapses and are questionable in the solar system microscopic physics [12] lead to conclude that the proper avenue towards unifying these two theories and thereby explaining the cosmological evolution completely will be understanding the space-time singularity and extending the conventional quantum theory as the position-momentum Canonical commutation relations are in accordance with the corresponding minimum uncertainty relations [13].

VI. APPENDIX:A

We can obtain the one-form $(d\phi)_a$ from a zero-form (a scalar field) ϕ in an explicit coordinate variables notation:

$$(d\phi)_a = \sum \frac{\partial \phi}{\partial x^\mu} [(dx)_a]^\mu \quad (21)$$

here the range of the summation is the dimension of space-time and it does not represent an infinitesimal change in ϕ . When expressed in a particular coordinate basis $[(dx)_a]^\mu$ will be just dx^μ , a coordinate one-form, and the μ -th component of $(d\phi)_a$ is $\frac{\partial \phi}{\partial x^\mu}$ as an arbitrary tensor T , in its operator form, represented in a coordinate basis can be expressed as:

$$\mathbf{T} = T^{\alpha\beta\dots}{}_{\mu\nu\dots} \left(\frac{\partial}{\partial x^\alpha}\right) \left(\frac{\partial}{\partial x^\beta}\right) \dots (dx^\mu)(dx^\nu) \dots \quad (22)$$

Here $(\frac{\partial}{\partial x^\alpha})$, $(\frac{\partial}{\partial x^\beta})$ are coordinate unit vectors and dx^μ, dx^ν are coordinate one-forms.

In terms of a coordinate basis the covariant d Alembertian operator can be obtained from the invariant:

$$(d\phi)^a (d\phi)_a = \sum \sum g^{\mu\nu} \frac{\partial \phi}{\partial x^\mu} \frac{\partial \phi}{\partial x^\nu} \quad (23)$$

here the explicit summations are again over μ, ν with the ranges same as above. When expressed explicitly in a coordinate basis the Lagrangian density of a massless scalar field is given by: $g_{\mu\nu} \partial^\mu \phi \partial^\nu \phi = g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi$. The infinitesimal change $\delta\phi$ of the scalar field ϕ can be interpreted as the scalar product of the one form $(d\phi)_a$ and the infinitesimal vector line elements.

VII. APPENDIX:B

Quantum field theory is the quantum theory of fields. It gives the dynamics of fields, the quantum probability amplitudes of creation and annihilation of particles, in contrast to quantum mechanics which gives the dynamics of the particles themselves obeying quantum principle. For the same boundary conditions these two descriptions match in the form of their kinematic solutions. Only for the free particle boundary condition the conventional interpretation of propagators in Q.F.T as giving the probability of particle propagation is in accordance with reality as the quantum probabilities are nowhere vanishing in both the theories. For microscopic particle physics experiments the free particle boundary condition is a good approximation in practice but ideally the field ϕ (or the quantum mechanical wave function ψ) is spatially confined within the experimental apparatus. In one dimension it is meaningless to say that a particle is propagating from one point to another if the probability of finding (or creation of) the particle is vanishing at some intermediate points.

For loops in one dimension, the momentum-space calculations give the probability that a pair of particles with given four-momentums are created at one space-time point and a pair of particles are annihilated at

another space-time point with the same four-momentums. This feature is transparent if one consider all possible space-time particle trajectories to form loops in one dimension which cannot be possible without the possible space-time particle-antiparticle (originated with given four-momentums and annihilated with the same four-momentums) trajectories crossing each other at least once in between any two given space-time points. Similar feature will be apparent if one interpret the loop as a particle encircling between any two given space-time points with the four-momentums at these two given space-time points remaining the same.

Let us illustrate this feature following the standard literature. We consider situations where free particle approximation hold. In vacuum at a given space-time point x (in one dimension), the particle- antiparticle pair production probabilities with two-momentums p_1, p_2 are $|\exp(-ip_1.x_1)|^2$ and $|\exp(-ip_2.x_2)|^2$ respectively apart from normalizing factors. Once produced the quantum mechanically allowed stationary state position-space wave functions that are available to the particle-antiparticle pairs are $\psi_P(x) = \exp(-ip_1.x_1)$ and $\psi_{AP}(x) = \exp(-ip_2.x_2)$ respectively where x denotes space-time points. The quantum mechanical joint probability that the particles produced at $x_1, x_2 = 0$ with two-momentums p_1 and p_2 can again coincide at a space-time point between x and $x + dx$ is:

$$P_x(p_1, p_2) = N_1 \frac{[dx]^2}{T^2 L^2} \quad (24)$$

here N_1 is the relative pair creation probability at the space-time point $x = 0$.

$P_x(p_1, p_2)$ is independent of $x, (p_1, p_2)$ and $\rightarrow 0$ as for free particle approximation $L \rightarrow \infty$ although the total probability of coincidence is unity when integrated over all space-time points. Hence quantum mechanically, numerous amount of particle-antiparticle pairs should be observed in any microscopic experiment performed during finite time-interval if there would have been spontaneous pair creations in vacuum.

In passing we note that a space-time formed out of loops in vacuum, closed time-like curves, as the source can not have an intrinsic time orientation in contrary to what is realized in nature.

We next note that in non-relativistic quantum mechanics the total joint probability that two distinguishable particles with energy E and momentums $(k, -k; k = \frac{2n\pi}{L}, |n| \gg 1)$ can coincide at some point x (here x is position only) is:

$$P(all) = \frac{4}{L^2} \int \int [\int \sin^2(kx_1) \delta(x_1 - x_2) \sin^2(kx_2) dx_2] [dx_1]^2 \quad (25)$$

here the integrals are performed over the interval $[-L/2, L/2]$. This expression turns out to be unphysical as the corresponding probability turns out to be unphysical [$P(all) = 3/2$]. Whereas classical mechanically the maximum value of $P(all)$ can be nearly unity when the particles suffer impulsive elastic collisions to stick together and come to rest at some point x_0 .

Similar features as discussed in this section in the contexts of equ.(24) and equ.(25) will appear in three dimensions.

In semiconductor physics, as charge carriers holes, are fictitious objects introduced for simplifications. In reality, quantum mechanically in a p-type semiconductor the motion of holes are out of the movements of the valence band or the acceptor level electrons. What is the proper explanation of the polarity of the Hall potential in a p-type semiconductor? The electrons experience forces (qvB) in the same direction as that of the holes and in terms of the motion of the electrons the polarity of the Hall voltage should be opposite.

VIII. APPENDIX:C

In this section we first consider the action for the gravitational field [20]:

$$S_g = -\frac{c^3}{16\pi k} \int G \sqrt{-g} d^4x \quad (26)$$

where G_{ab} is Einstein's tensor. General Theory of Relativity interwinds inertial mass (in general energy-momentum) of matter with space-time through the principle of equivalence and the dimension of the coupling constant k ($k = 6.67 \times 10^{-8} cm^3 - gm^{-1} - sec^{-2}$) is completely determined in terms of only mass, space and time unlike, for example, in electrodynamics where one have another fundamental quantity (electric charge) to determine the dimension of the coupling constant. This feature is transparent if we compare the

Newtonian-limit of the general theory of relativity with the Coulomb's law. The consequence of this feature is the following:

If we set $c, \frac{\hbar}{2\pi} = 1$ the dimension of k is length-squared ($[l^2]$) and it is no longer possible to set $k = 1$ as this will make the concept of space-time dimensions meaningless. Alternatively we could have set $c, k = 1$ (see footnote: page no. 269, [20]) then Planck's constant become dimensionful ($[l^2]$).

However we can set Boltzmann constant ($k' = 1$) by giving temperature the dimension of energy. In the reduced units $c, \frac{\hbar}{2\pi}, k' = 1$ the gravitational action becomes:

$$S_g = -K \int G \sqrt{-g} d^4x \quad (27)$$

where K have dimension $[l^{-2}]$.

We now conclude our discussion on Quantum Field Theory. In the rest of this section we will use the convention that $\langle \alpha | \alpha \rangle$ gives us probability density.

We first consider non-relativistic quantum mechanics in one dimension. In position-space the normalized quantum mechanical wave function ψ gives us the probability amplitude. $(\psi^* \psi) dx$ gives us the probability of finding the particle within the infinitesimal length interval dx . For a free particle one adopts the delta function normalization scheme for the quantum mechanical wave function:

$$\int_{-\infty}^{\infty} \psi_{k_1}^*(x) \psi_{k_2}(x) dx = \delta(k_1 - k_2) \quad (28)$$

In this equation the left-hand side is dimensionless while the one-dimensional delta function has dimension of length $[l]$ as is obvious from it's definition:

$$\int_{-\infty}^{\infty} f(k) \delta(k - k_0) dk = f(k_0) \quad (29)$$

for a regular function $f(k)$.

It would be appropriate to replace the free-particle boundary condition by periodic boundary condition which is a reasonable approximation in situations where free-particle boundary conditions hold as for a large length interval the spacing between the adjacent values of the momentum allowed by the periodic boundary condition is negligible.

In the reduced units ($c, \frac{\hbar}{2\pi} = 1$) the action is dimensionless. The action for a complex scalar field is given by:

$$I = \int L d^4x \quad (30)$$

where the covariant lagrangian density for a massive field is given by,

$$L = \frac{1}{2} \frac{\partial \phi^*}{\partial x_\mu} \frac{\partial \phi}{\partial x^\mu} - \frac{m^2}{2} \phi^{*2} \phi^2 \quad (31)$$

Consequently the dimension of ϕ (ϕ^*) should be inverse of length ($[l]^{-1}$). In the second quantization scheme $\langle \beta | \phi | \alpha \rangle$ replaces the classical field [21] and the expression $\langle \alpha | \phi^* \phi | \alpha \rangle$ gives the probability density of creation or annihilation of particles. For free-particle boundary conditions the Euclidean-space generating functional for a real scalar field is given by [22]:

$$W_E[J] = N_E e^{-S_E[\phi_0, J] - \frac{1}{2} Tr \ln [(-\bar{\partial}_\mu \bar{\partial}^\mu + m^2 + V''(\phi_0))_1 \delta(\bar{x}_1 - \bar{x}_2)]} \quad (32)$$

The terms in the logarithm giving quantum corrections are not dimensionless and the third term is not of the same dimension as of the first two terms.

For a real scalar field confined within a finite volume box with periodic boundary condition and consistent with the second quantization scheme we have (equ.3.28,[10]):

$$\phi(x) = \frac{1}{L^{3/2}} \sum_{(n_1, n_2, n_3)} \frac{\phi_{n_1, n_2, n_3}^+}{\sqrt{2k^0}} \exp\left[\frac{2\pi i}{L}(n_\mu x^\mu)\right] + \frac{1}{L^{3/2}} \sum_{(n_1, n_2, n_3)} \frac{\phi_{n_1, n_2, n_3}^-}{\sqrt{2k^0}} \exp\left[-\frac{2\pi i}{L}(n_\mu x^\mu)\right] \quad (33)$$

in a relativistic theory a covariant normalization using four volume would be appropriate and the normalizing factor should have dimension $[l^{-2}]$. Clearly the dimension of ϕ ($[l^{-\frac{3}{2}}]$) in equ.(33) is no longer $[l^{-1}]$ and equ.(31) no longer gives us a lagrangian density.

In other words the dimension of the scalar field ϕ as required from the action determining it's space-time evolution does not mach with the dimension required in the second quantization scheme in order that one can interpret $\langle \alpha | \phi^* \phi | \alpha \rangle$ as giving the probability density of creation or annihilation of particles. One can absorb the the normalizing factor into the the fock state by multiplying it by a factor with dimension $[l^{-1}]$ as in the second quantization scheme $\langle \beta | \phi | \alpha \rangle$ replaces the classical field. This will be in accordance with the probabilistic interpretation of the field ϕ as we have,

$$\langle \alpha | \phi^* \phi | \alpha \rangle = \langle \alpha | \phi^* | \beta \rangle \sum_{\beta} \langle \beta | \phi | \alpha \rangle . \quad (34)$$

where the sum is taken over all possible states.

However this will violate the interpretation of the Fock states as quantum mechanically the normalization of probability density $\langle \alpha | \alpha \rangle$ in a relativistic theory requires that the each of the normalizing factors for the Fock states, whose number depends on the number of particles present in the Fock state, should have dimension $[l^{-2}]$.

We now consider fermions and electromagnetic fields.

The covariant lagrangian density for each component of the free fermion fields (e.g, electrons-positrons) which is formed out of their causal space-time motion is given by:

$$L = \frac{1}{2} \frac{\partial \psi^*}{\partial x_{\mu}} \frac{\partial \psi}{\partial x^{\mu}} - \frac{m^2}{2} \psi^* \psi^2 \quad (35)$$

and dimension of ψ is again $([l]^{-1})$. After linearization of the second order partial differential equation satisfied by ψ we get the Dirac equation:

$$(i\gamma^{\mu} \partial_{\mu} - m)\psi = 0 \quad (36)$$

Hereafter the following lagrangian density is used to study Q.E.D:

$$L' = \bar{\psi}(i\gamma^{\mu} \partial_{\mu} - e\gamma^{\mu} A_{\mu} - m)\psi - \frac{1}{4} F_{\mu\nu}(x) F^{\mu\nu}(x) \quad (37)$$

with the dimensions of ψ as determined above the dimension of the first part of L' is no longer that of a Lagrangian density and the action formed out of it is not dimensionless in the reduced units. Also the current density, $e\bar{\psi}\gamma^{\mu}\psi$ (although the four divergence vanishes), do not involve any momentum operator and it is not obvious whether it is possible to have, in any approximation, the conventional interpretation of current density as charge-density times velocity from this expression.

The Lagrangian density of a charged scalar field which is similar to the quadratic Lagrangian density for Q.E.D is given by,

$$L_T = L_f + L_{em} + L_{int} \quad (38)$$

This complete Lagrangian density for a charged scalar is gauge invariant only if we take L_{int} to be,

$$L_{int} = -A_{\mu} j^{\mu} + e^2 A^2 \phi^* \phi \quad (39)$$

the second term do not have a transparent interpretation unless we consider screening effects from classical electrodynamics similar to the corresponding discussions given in Appendix:II of this article.

IX. APPENDIX: D

In this section we will illustrate the discussions in the context of equ.(20) in section IV. The metric of the two-sphere $S^2(\theta, \phi)$ is given by

$$ds^2 = \sqrt{A/4\pi}(d\theta^2 + \sin^2 \theta d\phi^2) \quad (40)$$

Here $0 \leq \theta \leq \pi$ and $0 \leq \phi \leq 2\pi$. For the unit two-sphere we have,

$$ds^2 = d\theta^2 + \sin^2 \theta d\phi^2 \quad (41)$$

with the ranges of θ, ϕ same as above. This coordinate system have the following ill-behavedneses:

- (i) The coordinate ϕ suffers a discontinuity along some direction from 2π to 0.
- (ii) ϕ is degenerate at the poles $\theta = 0, \pi$. In spherical polar coordinate system (r, θ, ϕ) the point $(r = c, \theta = 0)$ where c is a finite constant is obtained through identifying any two arbitrary points on a circle characterized only through distinct values of ϕ . Similar construction is valid for the point $(r = c, \theta = \pi)$. This will be obvious if we construct a two-sphere $S^2(\theta, \phi)$ from a two-dimensional circular strip by identifying the inner-boundary and the outer-boundary to two distinct points [see the discussions below eqn.(50) regarding the reduction of eqn.(48) to eqn.(50) for points on the polar axis]. This construction can be generalized to higher dimension.

- (iii) The metric is singular at the poles [see the above discussions and App.B (a point can't be obtained from a one-dimensional line-element without breaking the corresponding continuous topology)].

When the $S^2(\theta, \phi)$ can be embedded in the three dimensional Euclidean space one can introduce Carteian coordinate system (x, y, z) through the coordinate transformation:

$$x = \sin \theta \cos \phi, \quad y = \sin \theta \sin \phi, \quad z = \cos \theta. \quad (42)$$

here $x^2 + y^2 + z^2 = 1$. Although the metric is regular in the Carteian coordinate system the transformation coefficients $(\frac{\partial x'^\mu}{\partial x^\nu})$ are singular at the poles and also at some isolated points on the $x - y$ plane demonstrating the discussions below equ.(20). On the other hand to obtain the Spherical metric, singular at two isolated points, from the regular Carteian metric one has to introduce coordinate transformation with singular transformation coefficients.

We can also introduce two homeomorphic stereographic projections to coordinatize $S^2(\theta, \phi)$ embedded in R^3 . The first one is from the North pole $\theta = 0$ on the Equator plane to coordinatize the Southern hemisphere $\frac{\pi}{2} \leq \theta \leq \pi$. We have,

$$X = \cot \frac{\theta}{2} \cos \phi, \quad Y = \cot \frac{\theta}{2} \sin \phi \quad (43)$$

and this transformation is a homeomorphism at $\theta = \pi$. The second stereographic transformation is from the South pole $\theta = \pi$ on the Equator plane to coordinatize the Northern hemisphere $0 \leq \theta \leq \frac{\pi}{2}$. We have,

$$U = \tan \frac{\theta}{2} \cos \phi, \quad V = \tan \frac{\theta}{2} \sin \phi \quad (44)$$

and this transformation is a homeomorphism at $\theta = 0$.

The transformation between the (X, Y) and (U, V) coordinate systems is a diffeomorphism at their common domain $\theta = \frac{\pi}{2}$. The metric is also regular in these coordinate systems.

However the transformation coefficients between (X, Y) and (θ, ϕ) coordinates are singular at the South pole $(\frac{\partial X}{\partial \theta}, \frac{\partial Y}{\partial \theta} = 0 \text{ at } \theta = \pi)$, i.e, this transformation is not a diffeomorphism. Similarly the transformation $(\theta, \phi) \rightarrow (U, V)$ is not a diffeomorphism. It can also be shown that the transformations are not homeomorphism at the poles. Thus S^2 is not homeomorphic to the infinite plane (R^2). Also a geodesic (an element of a particular family of reparameterization invariant curves) on R^2 can never come close to intersect itself whereas geodesics on S^2 are closed curves.

We now consider the Robertson-Walker cosmological model. The space-time metric in terms of comoving isotropic observers is:

$$ds^2 = -d\tau^2 + a^2(\tau)[d\psi^2 + \sin^2\psi(d\theta^2 + \sin^2\theta d\phi^2)] \quad (45)$$

here $0 \leq \psi \leq \pi$, $0 \leq \theta \leq \pi$ and $0 \leq \phi \leq 2\pi$. The constant-time spatial three surfaces $\sum_\tau$ are compact (topologically S^3) and there is no four-dimensional spatial geometry available to embed $\sum_\tau$. The spatial metric is singular along the closed line elements $\theta = 0$, $\theta = \pi$ including the two point-poles $\psi = 0, \pi$.

The discussions in section IV [equ.(20)] and in this section show that metric singularities cannot be removed by diffeomorphically equivalent coordinate transformations. Thus the black hole and the cosmological metric singularities are unavoidable aspects of nature.

X. APPENDIX: E

In this section we consider the electrostatic potential of a polarized dielectric system. The electrostatic potential of a polarized dielectric system is given by,

$$V = \frac{1}{4\pi\epsilon_0} \int \frac{\vec{P} \cdot \vec{R}}{R^2} dv \quad (46)$$

here $\vec{P}$ is the polarization vector of the dielectric material and $\vec{R}$ is the vector joining the infinitesimal volume element dv carrying a dipole moment $\vec{P}dv$ to the point of observation. It's magnitude is given by eqns.(46). We can reduce equ.(55) to a simpler form consisting two terms: one from a bound surface charge density σ_b and another from a bound volume charge density ρ_b ,

$$V = \frac{1}{4\pi\epsilon_0} \int_{surface} \frac{\vec{P} \cdot \vec{da}}{R} - \frac{1}{4\pi\epsilon_0} \int_{volume} \frac{\vec{\nabla} \cdot \vec{P} dv}{R} \quad (47)$$

Here $\sigma_b = \vec{P} \cdot \hat{n}$, $\hat{n}$ is the normal to the surface of the material and $\rho_b = -\vec{\nabla} \cdot \vec{P}$.

The total volume charge density in presence of a polarized dielectric medium is given by:

$$\rho = \rho_b + \rho_f + \rho_{sb} \quad (48)$$

where we include $\sigma_b = \vec{P} \cdot \hat{n}$ in the free volume charge density as ρ_{sb} through the introduction of a proper delta function. For example in the case of a dielectric sphere we have,

$$\rho_{sb}(r, \theta, \phi) = \sigma_b(\theta, \phi) \delta(r - r_s) \quad (49)$$

Here r, r_s are radial distance (not vectors) and the delta function have dimension inverse of length. We then have:

$$\epsilon_0 \vec{\nabla} \cdot \vec{E} = \rho = \rho_b + \rho'_f \quad (50)$$

where $\rho'_f = \rho_f + \rho_{sb}$ and the divergence of the electric displacement vector $\vec{D}$ is given by,

$$\vec{\nabla} \cdot \vec{D} = \rho'_f \quad (51)$$

Everywhere apart from the surface of the dielectric we have $\rho'_f = \rho_f$ and the above equation (51) matches with the conventional expression for the divergence of $\vec{D}$:

$$\vec{\nabla} \cdot \vec{D} = \rho_f. \quad (52)$$

The effect of ρ_{sb} should be taken into the boundary conditions for $\vec{D}$. This will also have consequences to obtain the energy density of a given electrostatic configuration in presence of dielectric mediums [23] as,

$$\vec{\nabla} \cdot \vec{D} - \vec{\nabla} \cdot \vec{D}_0 = \rho_{sb} \quad (53)$$

We conclude this section with a few comments regarding the electrostatic field energy in presence of dielectrics.

The electrostatic field energy in presence of dielectric mediums can approximately be considered to consist of three parts [24]:

$$W_{tot} = \frac{1}{2} \int D \cdot E d\tau = \frac{\epsilon}{2} \int E \cdot E d\tau = W_{free} + W_{spring} + W_{bound} \quad (54)$$

here $\epsilon = \epsilon_0(1 + \chi_e)$. We briefly explain the three terms considering the realistic case of a dielectric filled charged parallel-plate capacitor:

i) W_{free} is the energy to charge the capacitor to produce the configuration with a given electric field. We can regain this energy if we discharge the capacitor by connecting the two plates through a conductor.

ii) W_{spring} is the energy required to increase the atomic/molecular dipole moments or to polarize the atoms/molecules depending on, respectively, whether the atoms/molecules have permanent dipole moments or not. This energy will be regained as heat when we discharge the capacitor.

iii) W_{bound} is the energy required to polarize the dielectric as a whole. The dipole-dipole interaction energy for two dipoles with dipole moments $\vec{p}_1$ and $\vec{p}_2$ and separated by $\vec{r}$ is:

$$U = \frac{1}{4\pi\epsilon_0} \frac{1}{r^3} [\vec{p}_1 \cdot \vec{p}_2 - 3(\vec{p}_1 \cdot \hat{r})(\vec{p}_2 \cdot \hat{r})] \quad (55)$$

U is minimum when the dipoles are antiparallel and maximum when the dipoles are parallel. Consequently for any statistically infinitesimal volume of the dielectric (i.e, volume elements which are very small compared to the dimension of system but large enough to contain sufficient number of atoms/molecules so that microscopic fluctuations can be approximately averaged to zero) the orientation of the atomic/molecular dipoles will be as isotropic as possible. To polarize the dielectric we have to orient the atomic/molecular dipoles in near-parallel configuration in a given direction and supply energy to increase the electrostatic energy of the dielectric. This energy, W_{bound} , will be regained as heat if we discharge the capacitor.

XI. SUPPLEMENT:I

We will now study the behavior of the covariant Klien-Gordon equation in the near horizon limit.

We will first consider the spectrum of the covariant Klien-Gordon equation in the $(3 + 1)$ -dimensional constant curvature black hole background which contains a one dimensional fixed point set of the time-like Killing vector field. This black hole space-time was obtained by Prof. M. Bannados, Prof. R. B. Mann and Prof. J. D. E. Creighton through the identification of points along the orbits of a discrete subgroup of the isometry group of the anti-de Sitter space-time. They used a static coordinate system where the constant-time foliations become degenerate along a particular direction apart from the black hole event horizon giving a one-dimensional fixed point set of the time-like Killing vector. The metric in the Schwarzschild like coordinates is given by,

$$ds^2 = \frac{l^4 f^2(r)}{r_h^2} [d\theta^2 - \sin^2 \theta (dt/l)^2] + \frac{dr^2}{f^2(r)} + r^2 d\phi^2 \quad (56)$$

where $f(r) = (\frac{r^2 - r_h^2}{l^2})^{\frac{1}{2}}$. These coordinates are valid outside the horizon ($r > r_h$) for $0 \leq \theta \leq \pi$ and $0 \leq \phi < 2\pi$. It is clear that the constant-time foliation becomes degenerate along the direction $\theta = 0$ and $\theta = \pi$ giving to a one-dimensional fixed point set of the time-like Killing vector field.

The covariant wave equation of a minimally coupled massive scalar field is given by,

$$\frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu \psi) - m^2 \psi = 0. \quad (57)$$

The solution of the angular equation is given by,

$$P_\nu^\mu(x) = \frac{1}{\Gamma(1-\mu)} \left(\frac{1+x}{1-x} \right)^{\frac{\mu}{2}} F(-\nu, \nu+1; 1-\mu; \frac{1-x}{2}) + c.c \quad (58)$$

where $x = \cos \theta$, $\mu = iEl$ and $(\nu + \mu) \neq$ an integer. Here $F(-\nu, \nu+1; 1-\mu; \frac{1-x}{2})$ is the hypergeometric function. This solution is C^1 throughout the angular range $0 \leq \theta \leq \pi$. Consequently the energy spectrum is continuous with divergent density of states.

We will now illustrate that the divergent density of states is a characteristic feature of the fixed point set of the time-like Killing vector field indicating the breakdown of the canonical formalism of the conventional quantum mechanics.

We will illustrate this feature in the context of the Schwarzschild black hole which contains a two dimensional fixed point set (the event horizon) of the time-like Killing vector field. Since the Hawking radiation through which the non-unitary black hole evaporation takes place originates mostly from the near horizon region we will consider the behavior of the spectrum of the covariant K-G equation in the near horizon region. Since the space-time foliation is static we will consider the stationary states. We will consider the radial solution of the covariant K-G equation. For the Schwarzschild black hole the constant-time foliations become degenerate at the black hole event horizon and it is not obvious to impose any consistent boundary condition on the horizon although the radial solution can be obtained through the WKB approximation and the spectrum is obtained using the semi-classical quantization condition (Brick wall model). However the conventional interpretation of the boundary condition is not obvious for extreme charged black holes.

To obtain the energy spectrum one impose the following boundary conditions on the wave function:

$$\psi = 0 \quad (59)$$

for $r = 2M + h$. The internal energy is given by [29]:

$$U = \frac{2\pi^3}{15h} \frac{2M^4}{\beta} \quad (60)$$

Similar behavior is also obtained when one considers the behaviour of matter fields in the Taub-NUT space-time which contains a zero dimensional fixed point (in the Euclidean sector) of the time like Killing vector field. In this case the angular solution (similar to spin-spherical harmonics) satisfies only the regularity condition that the angular part of probability current density integrated over S^2 is finite. This angular solution is similar to the spin-spherical harmonics.

The divergence of the density of states as $h \rightarrow 0$ as discussed above is a characteristic aspect of both the black hole event horizon and the cosmological event horizon.

We now make some comments regarding relativistic quantum mechanics similar to App:D. For relativistically covariant normalization of the quantum mechanical wave function (or each component of the wave function for spinors) of we have,

$$\int \psi^* \psi d^4x = 1 \quad (61)$$

This indicates that in the reduced units ($c, \frac{h}{2\pi} = 1$) the dimension of ψ is inverse length squared $[l^{-2}]$. While the lagrangian leading to the Klien-Gordon equation is given by:

$$L = \frac{1}{2} \frac{\partial \psi^*}{\partial x_\mu} \frac{\partial \psi}{\partial x^\mu} - \frac{m^2}{2} \psi^* \psi \quad (62)$$

The action determining the space-time evolution of ψ is dimensionless in the reduced units. This gives the dimension of ψ to be $[l^{-1}]$ in contradiction to that $[l^{-2}]$ obtained from the normalization condition.

In passing we note that to have a consistent time orientation for any space-time manifold, which through the principle of equivalence is form of existence of matter fields, particles should follow a particular family of reparameterization invariant curves in the space-time manifold [5].

XII. SUPPLEMENTII: A FEW COMMENTS ON DOUBLE SLIT INTERFERENCE

In this section we will give a causal description of double slit interference experiments. We will assume that we are working in the Cartesian coordinates in flat space-time where in the Lorentz gauge the electromagnetic potentials and the fields are conjugate harmonics and the fields obey the wave equations.

This analysis of interference patterns is based on the fact that the velocity of the causal part of electromagnetic waves is finite [Supp.III] and the corresponding wave-fronts travel with finite velocity. For a purely action-at-a-distance theory the two superposing fields are at the same state at any field point as they reach simultaneously from the two sources. Thus there would be no redistribution of intensity and we won't get interference fringe pattern. When the wave-fronts are traveling with finite velocity the wave fronts that simultaneously reach at a screen point from the slits should have originated from the two slits at different instants of time. We will find that the causal description of interference experiments together with the universality of interference fringe patterns naturally lead us to conclude that the velocity of light in vacuum is independent of the velocity of the source and the velocity of light is same in all directions. The causal interpretation is also useful to determine the velocity of light from the interference experiments. We will describe the corresponding situation with wave-groups. In particular we will give the mathematical conditions for maxima and minima for interference experiments with finite wave-groups. We will also discuss some conflicts between the physical description of the interference process for the wave-groups and the Fourier transform description of the wave-groups. These discussions are significant in the context of the foundational aspects of non-relativistic and relativistic quantum mechanics. In particular we find that a super-luminal velocity like the phase-velocity of the relativistic wave-function can determine a physical phenomenon like the corresponding interference fringe pattern which in this situation determine the probability density of finding the particle on the screen and hence the probable space-time motions of the corresponding particle passing through the slits.

We will follow the diagram [13F] of section 13.3 of [26] for the following discussions.

In Figure.13F [26] the wave fronts that simultaneously reach at the screen point P from the slits S_1 and S_2 should have originated from the two slits at different instants of time. The wavefront from S_2 should originate at time $\frac{S_2A}{c}$ earlier than the corresponding interfering wave front that has originated at S_1 . The phase difference is: $\delta = \omega\Delta t = \omega\frac{S_2A}{c} = \frac{2\pi}{\lambda}S_2A$ where ω and λ are the angular frequency and the wave-length of the light respectively, S_2A is the path difference between two interfering waves.

If we consider a field point 'P' at any instant of time t' and with positive values of X and Y coordinates $E_1(x, y, t)$ from the upper slit is interfering with E_2 from the lower slit that is similar to E_1 that had passed through 'P' at time Δt earlier. Thus the interfering waves from the slit S_1 and S_2 is given by:

$$E_1 = A \sin(\omega t - kx) \text{ and } E_2 = A \sin[\omega(t - \Delta t) - kx] = A \sin[\omega t - k(x + \delta x)]$$

These expressions are consistent with the fact that at $x = 0$, i.e, at the slits we are considering wavefronts E_1 at time t' and wavefronts E_2 at time $t' - \Delta t'$.

Hereafter we can continue the conventional procedure discussed in [37] to obtain the fringe pattern. Similarly, to reach a particular screen point P simultaneously the wavefronts from different parts of an aperture should have originated from wavefronts at the aperture at different instants of time differing in phases. This gives a causal description of the diffraction patterns. These discussions lead us to the following conclusions:

(i) We should note that, with the frequency of the light being constant, we have the following condition on Δt : $\omega\Delta t = \pi$ for the first interference minimum and similar will be the situation with other minima and maxima. Thus $\Delta t(s)$ remain to be the same for all experimental set-ups. The universality of the interference fringe patterns (the universality of the path differences) indicates that the velocity of light in vacuum is independent of the velocity of the source and the velocity of light is same in all directions. A similar analysis is given in [37] in the context of Michelson-Morley experiment.

(ii) The interference fringe pattern can be used to experimentally determine the velocity of light with a source with known angular frequency (ω) from the expression for the phase difference $\delta = \omega\frac{S_2A}{c}$.

We now consider the corresponding situation with wave-groups. When the coherence-length is much larger than the separation between the slits and the screen it is easy to extend the arguments for monochromatic-light to show that we should be using the phase-velocity to determine the interference pattern.

Now let us consider the situation when the separation between slits and the screen is large compared to the coherence-length or is of the order of the coherence-length. We assume that **the shape of the wave-groups originating from the two slits remain the same at any instant of time** although they are traversing in different directions. To illustrate, the wave-group can originate from atomic transitions and are traversing in vacuum. Let the wave-group from the slit S_1 , E_1 , has arrived at the field point 'P' at a time t . If we want to have the the first minimum at 'P' the wave-group from the second slit, E_2 , should be arriving at 'P' at a time Δt later such that E_1 at 'P' is ahead of E_2 in phase by an amount equals to π . In other

words the shape of E_1 and E_2 at 'P' at the instant $t + \Delta t$ should differ by half the wave-length with E_1 ahead of E_2 . The required time-interval Δt is $\frac{1}{v_p} \frac{\lambda}{2}$. Thus we can proceed similar to the earlier cases as we have for the phase-difference: $\omega \Delta t = \frac{\omega}{v_p} \Delta x = k \Delta x$ and we have the first minimum when $\Delta x = \frac{\lambda}{2}$. Similarly the second minimum occurs when E_1 ahead of E_2 by one and half wave-length from the beginning of E_1 . Here we have assumed that the lengths of the complete waves within the wave-group remain the same throughout the wave-group. If this is not the case the angular frequency will be varying along the wave-group and the mathematical calculations will have to be generalized accordingly for different orders of maxima and minima. The physical conditions for maxima and minima remain to be the same. For a weakly dispersive medium the above physical arguments regarding the path differences remain valid but in this case both the the angular frequency and the phase velocity is varying along the complete waves with different wave-lengths within the wave-group. The value of the wave-length λ in the above discussions should be taken at approximately at the time when they both reach the point 'P', i.e, $t + \Delta t$ and when the wave-group is weakly dispersing the wave-length may be defined through averaging over the group and the length of the wave-group should be defined through time averaging. Similar considerations are required when we consider the higher order minima. The above physical arguments for interference maxima and minima remain to be valid even when the wave-group is strongly dispersive. In terms of the group velocity the above physical arguments can be put in the following mathematical form which remains to be valid even for strongly dispersing mediums:

For positive values of Y -coordinates the n -th order interference is maxima determined by the following condition on the group velocity vectors of the two interfering wave-groups,

$$[(\vec{v}_{g1} - \vec{v}_{g2})t] \cdot (\frac{\vec{v}_{g1}}{v_{g1}}) = \sum \lambda_i(t)$$

In the above equation $t = 0$ when the center of the incident wave-group passes through the slits and i is running from one to n . Depending on the relative value of the length of the wave-group and the separation of the slits there can be a region where the interference pattern is governed by the phase-velocity. The wave-lengths λ_i are in general different and may depend on time for dispersive wave-groups. Here we only assumed that the dispersion does not depend on the direction of propagation. Similar discussions are valid for negative values of Y -coordinates with the suffices interchanged.

The above discussions are also in accordance with the significance of the coherent sources. We demonstrate this with the simplifying assumption that wave-group contains a few numbers of complete wave-lengths and we can define an average wave-length and the phase-velocities are not differing much from one another throughout the wave-group so that we can assign a single value to the angular frequency and hence to the time-period. This is the situation with the wave-groups discussed in the above paragraph. Let the coherence-length of the wave-group be L . The approximate number of average wave-lengths in the wave-groups at the point 'P' is $n = \frac{L}{v_g} \frac{1}{T}$. Here T is the average time-period of the wave-group at the point 'P' (we will also have to consider a time averaging over the duration of the wave-group at 'P' to define L and λ if the wave is weakly dispersing). Following the discussions in the preceding paragraph the interval of time through which the front of E_2 should differ from the front of E_1 in arriving at 'P' so that it just touches the end of E_1 at 'P' is $\tau = \frac{n\lambda}{v_p} = \frac{L}{v_g} \frac{1}{T} \frac{\lambda}{v_p}$. Here λ is the average wave-length of the wave-group at the point 'P' (we will again have to consider a time averaging over the duration of the wave-group at 'P' if the wave is weakly dispersing as is considered to define the average time-period T). For a wave-group which encompasses a few number of complete waves with the wave-lengths are not differing largely from each other we can take $v_p = \frac{\lambda}{T}$ and we have $\tau = \frac{L}{v_g}$ which is the coherence-time as is expected.

One should note that the the physical arguments presented in this section determine the interference fringe pattern for a wave-group even with very few numbers of complete wavelengths although the corresponding Fourier transform can include an wide range of monochromatic waves differing in wave-lengths and may also in respective phase velocities. If we consider the Fourier transform the interference fringe pattern is determined by the principle of superposition and the phase velocities of the monochromatic waves. We give a comparative study of these two descriptions for a wave-group with few numbers (of the order of ten) of same complete wavelengths propagating in a non-dispersive medium. We also assume that the amplitude remains the same or varies little over the complete wave-group. To have the first interference minima (there can be negligible residual intensity associated with the uncanceled half wave-lengths one at the beginning of E_1 and one at the end of E_2) as determined from the elementary physical considerations discussed in this article the wavelengths of the Fourier transform components should be mostly odd integral (including one) dividends of a basic wave-length which is same as the wave-length of the complete waves within the wave-groups. The spectrum gets even more restricted if we consider higher order minima.

On the other hand the Fourier transform of a sin function centered at the origin is given by the following expression:

$$a(k) = (\frac{2}{\alpha - k}) \sin(\frac{\alpha - k}{2})L - (\frac{2}{\alpha + k}) \sin(\frac{\alpha + k}{2})L$$

Here the wave group is given by $\sin(\alpha x)$ extending from $-(L/2)$ to $(L/2)$. Thus the mathematical analysis is not in accordance with the elementary physical description although the physical description is what happens in reality. For only very long wave-groups for which mathematically the phase-velocity can be given by a sharply peaked value these two descriptions agree with each other. Thus light wave groups of finite extent in vacuum are fundamental in nature similar to soliton solutions. We will illustrate the above aspects later.

For electromagnetic wave-group or other wave-groups like the ripple wave-groups the energy density is related with the value of the fields and the phase-velocity gives us the velocity with which energy is transmitted within the wave-group. When we consider matter-waves the probability density does not depend on the wave function but on the amplitude and phase-velocity within a wave-group does not have the corresponding interpretation as is mentioned above for electromagnetic wave-groups or ripple wave-groups. For matter wave-groups the particle flux is related to the particle velocity. However if the electron is described by a long wave-group, i.e, the electron is having a sharply defined momentum or the separation between the slits and the screen is small compared to the length of the wave-group the discussions at the beginning of this section regarding nearly monochromatic waves remain to be valid to describe the corresponding interference fringe pattern. In these situations matter-wave phase velocity determines a physical phenomenon like the corresponding interference fringe pattern as is illustrated at the beginning of this section. These discussions are significant for matter waves as the phase-velocities are different for the non-relativistic and the relativistic wave-functions. The phase velocity of the wave-function for a relativistic electron is $\frac{c^2}{v}$ [45]. In the case of Davisson-Germer experiment with relativistic electrons, if the separation between the sources and the screen is small compared to the length of the wave-groups the interference pattern is determined by the phase velocity which is greater than light and not by the group velocity as is conventionally referred [45] which is same as the velocity of the particle. The discussions made in point(i) in the context of the velocity of light are important as the angular frequencies are different for the non-relativistic and relativistic electron wave-functions (and consequently the corresponding time-intervals Δt for the first maximum will also be different). As discussed in the point(ii) we have, say for the first maximum, $\delta = \omega \frac{S_2 A}{v_p} (= 2\pi)$. Here $S_2 A$ is the experimentally determined path-difference, ω is the relativistic expression of the angular frequency of the electron wave-function. Thus we have a way to determine the super-luminal phase-velocity experimentally which determines a physical phenomenon like the interference pattern determining the probable space-time motion of the particle. The corresponding wave-lengths are also different and the experimental interference pattern should be in accordance with the relativistic expression of the matter-wave wave-length for the electrons. To illustrate these aspect, the condition for first interference maximum lead to the following relation between the relativistic phase velocity (v_{pr}) and the non-relativistic phase velocity (v_{pnr}) (**with the assumption that the path-difference remaining the same in both cases**): $\frac{v_{pr}}{v_{pnr}} = \frac{2}{\beta^2 \sqrt{1-\beta^2}}$.

Whereas De-Broglie hypothesis lead to the following relationship between the two phase velocities: $\frac{v_{pr}}{v_{pnr}} = \frac{2}{\beta^2}$. Here $\beta = \frac{v}{c}$ where v is the velocity of the electron. Thus the interference fringe pattern should be different with the non-relativistic and relativistic wave-functions. An experimental verification of the interference fringe pattern with a super-luminal phase-velocity provides a fundamental basis for the relativistic matter wave-function. The otherwise will indicate that second-quantization is a more consistent description of the relativistic dynamics of microscopic particles. Similar situations arise in the context of the collapse of the quantum mechanical wave-functions and the EPR paradox [46]. In the interference experiments the wave-function is completely delocalized between the different maxima prior to collapsing to the point where the electron is striking the screen. Thus the collapse of the wave-function is a non-local phenomenon. In EPR paradox the quantum mechanical wave-function predicts that the spin or polarization state of one of a pair of particles can change non-locally due to the corresponding change in the spin or polarization state of the other particle. The probability density of finding a particle at some point is determined by the corresponding matter wave-function signifying the physical reality of the wave-function. In this article we find that under certain circumstances the probability density and hence probable space-time motions of a relativistic particle is determined by the super-luminal phase-velocity of the corresponding relativistic wave-function.

To conclude, in this article we considered a causal description of interference which is different from Huygens's description. We found that the causal description of interference experiments indicates that the velocity of light in vacuum is independent of the velocity of the source and the velocity of light is same in all directions. The causal interpretation is also useful to determine the velocity of light from the interference experiments. We outlined the corresponding situation with wave-groups and have given the precise conditions for maxima and minima for interference experiments with finite wave-groups. We have also discussed some conflicts between the physical description of the interference process for the wave-groups and the Fourier transform description of the wave-groups. These discussions are important for the

foundational aspects of non-relativistic and relativistic quantum mechanics. We also find that a super-luminal velocity like the phase-velocity of the relativistic wave-function can determine a physical phenomenon like the corresponding interference fringe pattern determining the probability density of finding the corresponding particle in the region behind the slits and hence the probable space-time motions of the particle passing through the slits.

The above arguments indicates that the proper expression for fringe-shift in the Michelson-Morley experiments [26] is $2\omega d \frac{v^2}{c^2}$, where ω is the frequency of light. We can not follow the procedure followed in [26] to obtain the path-difference as the velocity of light, as is assumed, is different in different directions and as discussed above it is not proper to calculate the phase-difference between the interfering waves by multiplying the path-difference by $\frac{2\pi}{\lambda}$ directly. The fact that we can have the interference fringe system by allowing one photon to emerge from the source at each instant indicate that we have to introduce the corresponding electromagnetic wave description as fundamental and this description do not depend on the width of the slits apart from diffraction effects.

For double slit interference experiments with electrons, the wave fronts are the position-space wave functions ψ and are of the form $N e^{i(\vec{k} \cdot \vec{r} - \omega t)}$. The interfering states are the states of the electron at the slits at two different instants of time. Only for the central maximum the two states at the two slits are the same.

ψ is the complete microscopic description of the electrons in the electron -beam interference experiments. It is so as there is no underlying ensemble representation of the probabilistic interpretation of ψ . If we have to assume that, unlike the electromagnetic interference experiments, ψ is vanishing at the slits for slit width less than that of the diameter of the electrons then the finite volume property should be a fundamental aspect in the microscopic description of the electrons. In other words ψ gives a microscopic description of the electron upto a certain length. We can also illustrate this aspect with the following question:

What is the quantum mechanical description of a Radon atom in a rigid box when the distance of consecutive nodes and antinodes of ψ is equal to or less than the diameter of the atom?

XIII. SUPPLEMENTIII: COMMENTS ON CLASSICAL ELECTRODYNAMICS

In this Supplement we will make a few comments regarding the basic laws of Classical Electrodynamics.

In section:A we will consider the multipole expansion of the Cartezian coordinate inverse distance electrostatic potential for a regular and extended charge distribution. We will show that a proper binomial expansion in terms of the spherical polar coordinates of the Cartezian coordinate inverse distance electrostatic potential of a point charge not at the origin lead us to a multipole expansion in terms of the Legendre polynomials. The binomial expansion considered in this section is the most general and differs from the conventional expansion and gives well-defined result everywhere apart from two points. We will find that the potential expressed in terms of the Legendre polynomials is not well-defined at $\vec{r} = -\vec{r}'$ and differs from the corresponding value obtained from the $(\frac{1}{R})$ Coulomb potential. This leads us to conclude that the Cartezian coordinate $(\frac{1}{R})$ Coulomb law for a point charge not at the origin and the expansion, equation (83), in terms of the Legendre polynomials are the same object only in a restricted sense. This will also be apparent from the discussions in Section:B. Nevertheless following the experimental observations both in the static situations as well as in the dynamic situations we have adopted the solutions expressed in terms of the Legendre polynomials to describe the potentials in the spherical polar coordinates. The discussions in this article lead us to conclude that the force law in the Cartezian coordinates, given by the inverse square law, and the force law in the spherical polar coordinates, given in terms of the Legendre polynomials, are different. The potential expressed in terms of the Legendre polynomials remains to be well-defined everywhere for a regular source.

In section:B we will consider the Poisson's equation in the static situation. It is not possible to apply the integral version of the divergence theorem if the vector field is not well-behaved. We will demonstrate in this section that in the spherical polar coordinates an expression analogous to the integral version of the divergence theorem for a point charge in Electrostatics is valid when the point charge is at the origin.

We will discuss the derivation of Maxwell's equation from the force laws. We find that in flat space time it is in general not possible to derive the local Maxwell's equations in the familiar form from the inverse square force laws if we use arbitrary curvilinear coordinates. The equations for both the potentials and the fields contains non-local terms irrespective of the choice of gauge. This is the situation with the spherical polar coordinates in flat space-time. In the spherical polar coordinates (r, θ, ϕ) , the divergence of the electric field and the curl of the magnetic field contain non-local terms if the fields are expressed in terms of the corresponding inverse-square laws. Similar discussions are valid in the dynamical situations. In the dynamical situations the well-known Green's function, eqn.(120), does not satisfy both the inhomogeneous

and the homogeneous wave equations in terms of the spherical polar coordinates (r, θ, ϕ) . We will discuss this issue in Section:IV. We will in particular consider the fields of an oscillating electric dipole. We will find that both the potentials and the fields obtained from the Green's function (120) do not obey the wave equations for points outside the sources and do not contain $(\frac{1}{r})$ radiation fields observed in the experiments. We will also find that the corresponding potentials violate the Lorentz gauge condition. This may also remain valid in curved space-time.

In this article we give a formulation of electrodynamics in an arbitrary curvilinear coordinates for arbitrary sources with the potentials as fundamental variables on the basis of the following assumptions:

Following the experimental observations, we adopt the principle of covariance to write the equations for the potentials in curvilinear coordinates and in curved space-time with reference to the physical laws that photons propagates along null geodesics in vacuum.

The potentials satisfy the homogeneous wave equations in vacuum and the modal solutions of the wave equations creates or annihilates the photons in a second quantization scheme in the curvilinear coordinate system.

The electromagnetic fields are related to the potentials by the conventional relations. However the electromagnetic fields need not to satisfy the wave equations in vacuum. In the Cartesian coordinates both the potentials and the fields obey the wave equations in vacuum. This is because of the fact that in terms of the flat space-time Cartesian coordinates the partial derivatives in the curl and in the wave equation commute. This may not be the situation in an arbitrary curvilinear coordinates where the corresponding partial derivatives may not commute. Mathematically the above discussions are related to the existence of conjugate pair of harmonics in an arbitrary curvilinear coordinate system in four dimensions. In the Cartesian coordinates where conjugate pair of harmonics exist the potentials satisfying the wave equations obey the Lorentz gauge and the fields also satisfy the wave equations in vacuum. This may not be the situation for the potentials satisfying the wave equations in an arbitrary curvilinear coordinate system where conjugate pair of harmonics may not exist and the corresponding electromagnetic fields may not satisfy the wave equations. This is the situation with the spherical polar coordinates in flat space-time. In the spherical polar coordinates only the divergence of $\vec{B}$ and Faraday's law remain to be valid for time dependent sources. These laws define the fields in terms of the potentials and should remain valid in a potential formulation of electrodynamics irrespective of the nature of the sources. The electromagnetic fields give the forces acting on a charged particle. The homogeneous wave equations for the potentials reduce to the corresponding Laplace's equations in the static limit of static sources and $c \rightarrow \infty$ and the corresponding solutions expressed in terms of the Legendre polynomials describe the experimentally observed static potentials. We will later demonstrate that in terms of an arbitrary curvilinear coordinate system it is in general not possible to find a gauge so that the potentials satisfy both the wave equations and the Lorentz gauge condition.

For static sources all the Maxwell's equations remain valid for points outside the source both in the Cartesian coordinates as well as in the spherical polar coordinates. In the static limit of time-independent sources and $c \rightarrow \infty$ the wave-equations for the potentials reduce to the corresponding homogeneous Laplace's equations. The current density should be steady, time independent and satisfy the corresponding regularity conditions discussed below equation (86).

We now consider the expressions for the potentials within the source for both the static as well as the dynamic situations. In some situations the potentials within the source are given by the inhomogeneous equations. This is the situation in the flat space-time with the Cartesian coordinates and the fields are given by the inverse square laws. On the other hand in some situations for points both inside and outside the sources the potentials are obtained from the homogeneous equations in terms of the multipole coefficients. The homogeneous equations do not contain the source charge-current densities. The relationship between the source and the fields are expressed by the multipole coefficients. The multipole coefficients are determined either from the assumed nature of the source or from the boundary conditions. The boundaries, the boundary conditions, permittivity and permeability will be different for points inside and outside the source. This is the situation with the spherical polar coordinates in flat space-time and the potentials are given by multipole expansions in terms of the Legendre polynomials. We will discuss the solutions in eqns. (83,84,127) which agree with the experimental observations and satisfy the homogeneous equations. Mathematically, the distributions for the potentials in the Cartesian coordinates are either the Green's functions of the Poisson's equation in the static situations or the Greens functions of the four dimensional wave equations in the dynamical situations. On the other hand in the if we use the spherical polar coordinates, taking divergence of $\frac{\vec{R}}{R^2}$ w.r.t $\vec{r}$ does not give us the same result if we take the divergence w.r.t $\vec{R}$ or $\vec{r}$ (with a negative sign). This is clear from eqn.(73) and (78) and in terms of the spherical polar coordinates (r, θ, ϕ) the inverse-distance potential of a point charge not at the origin is not the Green's function of the Poisson's equation. This also follows from the discussions in Section:A where we have proved that $\frac{1}{R}$ and the corresponding expansion in terms of the Legendre polynomials, eqn.(83), do not agree with each other for all points. In

this article, in the spherical polar coordinates, the distributions relating the potentials to the source for both the static and the dynamic cases are constructed out of the modal solutions of the homogeneous Laplace's equation or the four dimensional homogeneous wave equations respectively. Only for the particular case of a static point charge at the origin the electric field satisfies Poisson's equation in (r, θ, ϕ) with the charge density given by a delta function at the origin and the field is given by the monopole term. For Maxwell's equations in curved space-time, the discussions in this article demonstrate that the diffeomorphism invariance of the inhomogeneous Maxwell's equations may become a non-trivial issue even if one have the inhomogeneous Maxwell's equations in a particular curved space-time. Similar as above, photons travel along null geodesics in vacuum is a more fundamental principle in this respect.

The discussions in this article are significant for the corresponding Lagrangian formulations and quantum field theory. In terms of the spherical polar coordinates the lagrangian does not contain the interaction term explicitly. In a second quantization scheme solutions of the form eqns. (127,128,129..) are similar to the interaction action in the Cartezian coordinates and gives the coupling between the source and the electromagnetic field. The stress tensor for the electromagnetic fields in the spherical polar coordinates is to be obtained using the Lagrangian formulation. We discuss this issue for time dependent sources in Section:IV. The stress tensor in this situation is formulated in terms of the potentials.

We will conclude Section:B with a realistic description of the electrostatic behavior of conductors with reference to departure from the inverse square law. We will also improve the proof of the curl of $\vec{B}$ law and the no-work law for the electric field. We have also discussed the forms of the Maxwell's equations in the spherical polar coordinates, (r, θ, ϕ) , with sources depending explicitly on the fields like sources obeying the Ohm's law in this section.

In Supplement:X we will discuss the electromagnetic potentials in the spherical polar coordinates for time dependent sources. In this article we only consider accelerated charges. We will construct the solution for the potentials following the static situations. The potentials satisfy the four dimensional homogeneous wave equation in the spherical polar coordinates. For points outside the source the solutions are given by a set of new polynomials. We will give a few polynomials for field points outside the source. We will in particular consider the potentials and the fields due to an oscillating electric dipole. To the leading order the electromagnetic fields agree with the conventional expressions although the power radiated differ from the conventional expressions [24] through a different dependence on the polar angle. For an oscillating point charge the conventional Green's function in the spherical polar coordinates, equation (120), does not give $(\frac{1}{r})$ radiation fields in contrary to the experiments. We will also give the solutions of the homogeneous wave equation for the potentials for $r < r'$, where r is the field coordinate and r' is the source coordinate. The solutions reduce to the well-known multipole potentials expressed in terms of the Legendre polynomials in the static limit. The expression of the time-averaged power radiated from an oscillating electric dipole as is obtained in this article, eqn.(140), agrees with the observations.

A. Multipole expansion and the Legendre polynomials

In this section we will consider the binomial expansion of the $(\frac{1}{R})$ Coulomb potential in terms of the spherical polar coordinates (r, θ, ϕ) in a way different from the conventional procedure. We will find that this binomial expansion of the Cartezian coordinate $(\frac{1}{R})$ potential of a point charge not at the origin in terms of the spherical polar coordinates gives well-defined result for all points apart from two points.

We consider the electrostatic potential of an extended but localized charged distribution. We will consider points outside the source. Let $\vec{r}'(r', \theta', \phi')$ be the position vector for an infinitesimal volume element dv' within the source which makes an angle θ' with the positive Z polar axis and an azimuthal angle ϕ' w.r.t the positive X axis. Let $\vec{r}(r, \theta, \phi)$ be the position vector of the point of observation (P) making an angle θ with the polar axis and an angle ϕ with the positive X axis. The magnitude of the position vector $\vec{R}$ between dv' and P is then given by:

$$R^2 = r^2 + r'^2 - 2rr'\cos(\gamma) \quad (63)$$

where,

$$\cos(\gamma) = \cos\theta'\cos\theta + \sin\theta'\sin\theta\cos(\phi - \phi') \quad (64)$$

The electrostatic potential at P is given by,

$$V(P) = \frac{1}{4\pi\epsilon_0} \int \frac{\rho(r', \theta', \phi') r'^2 \sin \theta' dr' d\theta' d\phi'}{[r^2 + r'^2 - 2rr' \cos(\gamma)]^{1/2}} \quad (65)$$

In the conventional procedures [24] one expands $(1+z)^{-\frac{1}{2}}$ in terms of z where $z = x^2 - 2x\cos\gamma$ and $x = \frac{r'}{r}$. If we consider points near the South pole (w.r.t the field point), the conventional binomial expansion is not valid in general for all r' . The absolute value of the ratio of two successive terms is $\frac{n+1/2}{n+1}z$ and the maximum value of z is 3.

We now consider an alternate binomial expansion for the potential which is valid for all $r' > r$ and lead to the Legendre polynomials. For any point outside the source we have the following expression for the potential:

$$V(P) = \frac{1}{4\pi\epsilon_0} \int \frac{\rho(r') r'^2 \sin \theta dr' d\theta' d\phi'}{(r^2 + r'^2)^{1/2} [1 - 2\frac{rr'}{(r^2 + r'^2)} \cos(\gamma)]^{1/2}} \quad (66)$$

We first expand $[1 - 2\frac{rr'}{(r^2 + r'^2)} \cos(\gamma)]^{-1/2}$ in terms of $2\frac{rr'}{(r^2 + r'^2)} \cos(\gamma)$. We obtain the following expression for the potential:

$$V(P) = \frac{1}{4\pi\epsilon_0} \int \rho(r') r'^2 \sin \theta dr' d\theta' d\phi' \frac{1}{(r^2 + r'^2)^{1/2}} [1 + \frac{1}{2} \frac{2rr'}{r^2 + r'^2} \cos(\gamma) + \frac{3}{8} (\frac{2rr'}{r^2 + r'^2})^2 \cos(\gamma)^2 + \dots] \quad (67)$$

We thereafter multiply the resultant series by $(r^2 + r'^2)^{-1/2}$ and lastly we expand different powers of $(r^2 + r'^2)$ in terms of $\frac{r'}{r}$. For points outside the source $r > r'$. This expansion is valid for all points $r' \leq r$ apart from $\vec{r} = \pm \vec{r}'$. For $\vec{r} = \vec{r}'$ the potential is diverging and the corresponding binomial expansion may not remain well-defined. We will later consider the problem associated with $\vec{r} = -\vec{r}'$.

The resulting expression gives us a series in terms of the Legendre polynomials:

$$V(P) = \frac{1}{4\pi\epsilon_0} \int \frac{r'^l}{r^{l+1}} P_l(\cos \gamma) \rho(\vec{r}') dv' \quad (68)$$

Hereafter we can use the composition rule [23] for the Legendre polynomials to express the potential in terms of the Legendre polynomials in the field coordinates.

We note that for $\vec{r} = -\vec{r}'$ the above expansion is not well-defined. This is because the expansion of $\frac{1}{(1+x)}$ is not well-defined for $x = 1$. To illustrate, let us consider the potential of a point charge, q , situated at $z = a$. The potential expanded in terms of Legendre polynomials for $r > a$ is given by:

$$V(\vec{r}) = \frac{q}{4\pi\epsilon_0} \sum \frac{a^l}{r^{l+1}} P_l(\cos \theta) \quad (69)$$

For $z = -a$ the series in terms of the Legendre polynomials is indeterminate as is expected from the Leibnitz test [46]. It gives either zero or

$$V(\vec{r}) = \frac{q}{4\pi\epsilon_0 a} \quad (70)$$

Whereas Coulomb's law give us the following expression for electrostatic potential at $z = -a$:

$$V(\vec{r}) = \frac{q}{8\pi\epsilon_0 a} \quad (71)$$

Clearly the potential obtained at $z = -a$ from expansion in terms of Legendre polynomials is not well-defined and is not in accordance with Coulomb's law expressed in terms of the Cartesian coordinates. This is related with the fact that the expansion of $\frac{1}{1+z}$ is not well-defined for $z = 1$ and equations (70,71) show that $\frac{1}{1+z}$ and the corresponding binomial expansion may not be the same object. $(\frac{1}{R})$ is not the generating

function for the expression (83) for all points. We will later find that the inverse square law and, in general, the expansions in terms of the Legendre polynomials are solutions of different differential equations in different coordinate systems and it is again expected that they may not be the same object without restrictions. In terms of the spherical polar coordinates (r, θ, ϕ) the inverse square law and the corresponding expansion in terms of the Legendre polynomials satisfy different differential equations. We will find in Section:B that in the spherical polar coordinates (r, θ, ϕ) we can not interchange $\vec{r}$ with $\vec{R}$ or with $\vec{r}'$ (at the cost of a minus sign) while taking the divergence of $(\frac{\hat{R}}{R^2})$ and hence in the spherical polar coordinates (r, θ, ϕ) the inverse square law does not satisfy the Green's function equation for a point charge not at the origin. On the other hand with equation (83), for both $r > r'$ and $r' > r$, we can interchange $\vec{r}$ with $\vec{r}'$ while taking the divergence of the corresponding electric field. Both the divergences are vanishing and equation(83), the potential for a point charge not at the origin and expressed in terms of the Legendre polynomials, also does not satisfy the Green's function equation for a point charge not at the origin and is a solution of the homogeneous Laplace's equation. Only the $(\frac{1}{r})$ Coulomb potential for a point charge at the origin satisfy the corresponding Green's function equation in terms of the spherical polar coordinates (r, θ, ϕ) . We will discuss this issue in detail in Section:B.

Following the experimental observations we continue to describe the potentials in the spherical polar coordinates in terms of the Legendre polynomials irrespective of the fact that in general these expressions satisfy the homogeneous Laplace's equation in terms of (r, θ, ϕ) . Also in the dynamic situation the potential for points outside the source is given by equ.(127), which in a second quantization scheme is consistent with the experimental fact that photons travel along the null geodesics in vacuum, and expression (127) reduces to the corresponding part of $r > r'$ from eqn.(21) in the static limit of static source and $c \rightarrow \infty$. For a point charge not at the origin the potential expressed in terms of the Legendre polynomials is not well-defined at $\vec{r} = \vec{r}'$ and $\vec{r} = -\vec{r}'$. However for a non-singular volume charge density for which the potential $\frac{1}{4\pi\epsilon_0} \int \frac{r'^l}{r^{l+1}} P_l(\cos \gamma) \rho(\vec{r}') dv'$ and the corresponding electric field vanishes in the limit the source volume $\delta v \rightarrow 0$, the potential and the field expressed in terms of the Legendre polynomials remains finite everywhere and the field is also well-behaved at all points. We will discuss this issue in detail below equation (85).

B. Discussions on Poisson's equation

In this section we will consider the Poisson's equation for the electric field. We first note that we can not apply the integral version of the divergence theorem if the vector field is divergent at any point inside the region of interest. The expression $E_x(\Delta x) = E_x(r=0) + \Delta x [\frac{dE_x}{dx}]_{r=0}$ is not defined for a point charge at the origin. In one dimension we can use the relation: $\int_{x_1}^{x_2} (df/dx) dx = f(x_2) - f(x_1)$ only if $f(x)$ is well-behaved for all x between and including the limits. In the following section we will show that in the spherical polar coordinates an explicit volume integration of the divergence of the electric field of a point charge gives us the total charge when the charge is at the origin. Also for a point charge at the origin the integral of the divergence of the electric field is vanishing when the volume of integration does not include the origin. This together with the fact that the total flux of the electric field of a point charge at the origin over an S^2 centered at the origin is given by $\frac{q}{\epsilon_0}$ lead us to us a relation for the electric field of a point charge at the origin analogous to the integral version of the divergence theorem.

We will shortly find that when the point charge is not at the origin the electric field in the spherical polar coordinates (r, θ, ϕ) satisfying local differential equations is not given by the inverse square law. However in the following discussions we will ultimately consider a point charge at the origin and we use the inverse square law in (r, θ, ϕ) for the electric field of a point charge at the position vector $\vec{a}$. We integrate the divergence of the corresponding electric field over a sphere centered at the origin. The radius vector of the point charge is $\vec{a}$ and the radius of the sphere is greater than $|\vec{a}|$. The integral of the angular derivatives vanishes. For example the θ -integral of the polar angular derivative term is given by,

$$4\pi\epsilon_0 I_\theta = \int_0^\pi d\theta \frac{\partial}{\partial \theta} \sin \theta \int \frac{\rho(\vec{r}' - \vec{a})}{R^3} (\hat{\theta} \cdot \vec{r}') dv' \quad (72)$$

and this vanishes. Similarly the ϕ -integral of the azimuthal derivative term vanishes out of uniqueness. Here one should first integrate the source integral which contains a delta function at $\vec{r}' = \vec{a}$.

The r -integral of the radial derivative term gives us the following result,

$$4\pi\epsilon_0 I_r = \int d\Omega \frac{r^3 q}{[r^2 + a^2 - 2racos\gamma]^{3/2}} - \int d\Omega \frac{r^3 q(\vec{a} \cdot \hat{r})}{[r^2 + a^2 - 2racos\gamma]^{3/2}} \quad (73)$$

In the limit $a = 0$ this gives the result $I_r = \frac{q}{\epsilon_0}$. Hence we have $\int \vec{\nabla}_{\vec{r}} \cdot \vec{E} dv = \frac{q}{\epsilon_0}$ when the point charge is at the origin.

When the point charge is at the origin $4\pi\epsilon_0 \int \vec{\nabla}_{\vec{r}} \cdot \vec{E} dv$ is vanishing if the integrating volume does not include the origin. If we consider any annular region centered at the point charge at the origin, the volume integral of the radial derivative term of $\vec{\nabla}_{\vec{r}} \cdot \vec{E}$ is given by the following expression:

$$\int q d\Omega \left[\left(\frac{r_2}{r_1} \right)^3 - \left(\frac{r_1}{r_2} \right)^3 \right] = 0 \quad (74)$$

Whereas the θ -integral of the polar angular derivative term is,

$$\int_0^\pi d\theta \frac{\partial}{\partial \theta} \sin \theta \int \frac{\rho(r')}{R^3} (\hat{\theta} \cdot \vec{r}') dv' = 0 \quad (75)$$

we also have $\vec{r}' = 0$. The azimuthal part vanishes out of uniqueness. It is easy to show that the volume integral of $\vec{\nabla}_{\vec{r}} \cdot \vec{E}$ is vanishing for any arbitrary volume not including the point charge at the origin.

Thus we have the well-known Poisson's equation for a point charge at the origin,

$$\vec{\nabla}_{\vec{r}} \cdot \left(\frac{\hat{r}}{r^2} \right) = 4\pi\delta^3(\vec{r}) \quad (76)$$

The above discussions together with the fact that the total flux of the electric field over a closed surface is $\frac{Q}{\epsilon_0}$ allows us to write the corresponding integral version of Gauss's divergence theorem for the electric field of a point charge at the origin.

We can derive a corresponding expression when the charge is not at the origin following the above procedure through replacing $\vec{r}$ and $\vec{r}'$ by:

$$(\vec{R}, \vec{R}') = (\vec{r} - \vec{r}', \vec{r}'' - \vec{r}') \quad (77)$$

respectively and considering the explicit integrals as above over a sphere centered at $\vec{r}'$. The source integral is in terms of $\vec{r}'' - \vec{r}'$.

Thus we have the following relation,

$$\vec{\nabla}_{\vec{R}} \cdot \left(\frac{\hat{R}}{R^2} \right) = 4\pi\delta^3(\vec{R}) \quad (78)$$

where $\vec{R} = \vec{r} - \vec{r}'$.

We will later discuss that in general taking divergence with respect to the Carteizian coordinates do not give us the same result if we take the divergence with respect to the spherical polar coordinates. However the two coordinate systems give us the same result for the vector field $(\frac{\hat{r}}{r^2})$. The divergence in both the coordinates are zero although there is a source at the origin. Thus we can use eqn(76) and eqn(78) in the Carteizian coordinates also. The experimental verifications of flat space-time Quantum Electrodynamics in the Carteizian coordinates also lead us to these relations.

We should note that if we use the spherical polar coordinates, taking divergence w.r.t $\vec{r}$ does not give us the same result if we take the divergence w.r.t $\vec{R}$ or $\vec{r}'$ (with a negative sign). This is clear from eqn.(73) and (78). It is a unique feature of flat space-time metric expressed in terms of the Carteizian coordinates that we can interchange $\vec{r}$ and $\vec{r}'$ (at the cost of a negative sign) while taking the divergence of $\frac{\vec{R}}{R^3}$. The metric and hence R and the scale factors within the divergence operator are not symmetric w.r.t the field coordinates and the source coordinates if we use, say, the spherical polar coordinates. To illustrate the divergence of $\frac{\vec{R}}{R^3}$ w.r.t $\vec{r}$ when $\vec{r}$ and $\vec{r}'$ are collinear is given by $-\frac{2r'}{r(r-r')^3}$. The divergence of the same vector field w.r.t $\vec{r}'$ is

given by $\frac{2r}{r'(r-r')^3}$. It is obvious that the above two divergences are not antisymmetric. In fact in terms of the spherical polar coordinates (r, θ, ϕ) , the Poisson's equation for the inverse square field takes the following form:

$$\vec{\nabla}_{\vec{r}} \cdot \vec{E} = t_r + t_\theta + t_\phi \quad (79)$$

Here,

$$t_r = \int \frac{\rho(\vec{r}')}{r^2} \frac{\partial}{\partial r} \left[\frac{r^2(\vec{r} - \vec{r}') \cdot \hat{r}}{R^3} \right] dv' \quad (80)$$

and

$$t_\theta = \int \frac{\rho(\vec{r}')}{r \sin \theta} \frac{\partial}{\partial \theta} \left[\frac{(\vec{r}') \cdot \hat{\theta} \sin \theta}{R^3} \right] dv' \quad (81)$$

and

$$t_\phi = \int \frac{\rho(\vec{r}')}{r} \frac{\partial}{\partial \phi} \left[\frac{(\vec{r}') \cdot \hat{\phi}}{R^3} \right] dv' \quad (82)$$

For a non-singular volume charge density the terms t_r, t_θ, t_ϕ are in general logarithmically divergent when the field point is within the source and give non-local terms when the field point is outside the source. Also it is not possible to derive the other Maxwell's equation from the inverse square force laws and using the spherical polar coordinates. The equations for both the potentials and the fields contain strong non-local terms both in the static as well as in the dynamic situations. This is contradictory in a second quantization scheme where the modal solutions of the potential equations, as operators, are expected to create photons traveling in null geodesics in vacuum. On the other hand it is very well verified by experiments that in the static situations the expansions in terms of the Legendre polynomials describes the fields expressed in terms of the spherical polar coordinates for both points inside and outside the source. The multipole expansions in terms of the Legendre polynomials (more generally the associated Legendre polynomials) in general satisfy the homogeneous Laplace's equation in the spherical polar coordinates (r, θ, ϕ) . Only for a point charge at the origin the potential satisfy the Poisson's equation. As we have discussed earlier the field of a point charge not at the origin is not given by the inverse square law in (r, θ, ϕ) . The field can be found from the assumption that the potential satisfies Laplace's equation, from the fact that the potential is symmetric under the interchange of the field coordinates and the source coordinates (it can be easily seen from the following expression for the potential that the electric field is antisymmetric under the interchange of the source point and the field point when both of them are on the positive Z -axis and this relation is valid in general for the following expression as long as it remains well-defined) and from the boundary conditions at infinity. We first consider the simple situation of the azimuthally symmetric field of a point charge on the polar axis. The potential satisfying the above conditions are given by:

$$V(\vec{r}, \vec{r}') = \frac{q}{4\pi\epsilon_0} \sum \frac{r_{<}^l}{r_{>}^{l+1}} P_l(\cos \theta) \quad (83)$$

In the more general situations where the point charge is not on the polar axis $\cos \theta$ is replaced by $\cos \gamma$. Only for the particular case of a point charge at the origin the potential is given only by the monopole term and satisfies the Poisson's equation. We can obtain the field for an extended source by generalizing the above expression. For a regular source the field inside the source is given by the following expression:

$$4\pi\epsilon \vec{E}(\vec{r}) = -\vec{\nabla}_{\vec{r}} \int_{r < r'} \sum \left[\frac{r^l}{r'^{l+1}} \rho P_l(\cos \gamma) \right] - \vec{\nabla}_{\vec{r}} \int_{r \geq r'} \sum \left[\frac{r'^l}{r^{l+1}} \rho P_l(\cos \gamma) \right] \quad (84)$$

When the charge density is non-singular the field is well-defined everywhere inside the source. The divergence and curl of $\vec{E}$ is zero. Thus we find that in terms of the spherical polar coordinates, for points both

inside and outside the source, the static potential and the field is given in terms of the Legendre polynomials which satisfy the homogeneous equation, the Laplace's equation. One should note that if physically the above solution gives us the correct field, then through Helmholtz's theorem, in terms of the spherical polar coordinates, the Laplace's equation will be the correct equation for the electric field. The relationship between the solution and the source is obtained from the multipole coefficients defined over the complete source. In day-to-day problems one usually specifies the source and the multipole coefficients are obtained from the assumed form of the source density. In a more general situations the multipole coefficients are to be obtained from the boundary conditions. The boundaries, boundary conditions, permittivity and permeability will be different for points inside and outside the source. At the origin $\hat{r}, \hat{\theta}, \hat{\phi}$ are vanishing and degenerate. For a regular charge distribution one can find the field by studying the behavior of the field at points near the origin. Inside the source the velocity of light will be determined by the permittivity and permeability of the medium. We should note that in the spherical polar coordinates the divergence of the electric field is vanishing although there exist a monopole term in the electric field with the total charge dependent on the field coordinates within the source. The multipole coefficients are determined by integrating over the source coordinates and while taking the derivatives of the fields the multipole coefficients should be considered as constants. The divergence of the monopole field give us a delta function at the origin and hence is vanishing at the field point when the field point is not at the origin. If the charge density is regular the divergence is vanishing even if the field point is at the origin as $\rho\delta v \rightarrow 0$ in the limit $\delta v \rightarrow 0$. Only when we have a point charge at the origin the divergence of the electric field is non-vanishing which is not the situation with a regular charge distribution. However the monopole term in eqn.(84) gives finite flux for a spherical surface with radius r and centered at the origin. This is consistent with the integral version of Gauss's law if we chose the fields and the closed surfaces with some restrictions. For a sphere lying completely within the source and centered at the origin the field throughout the region should be given by the multipole coefficients evaluated at the boundary two-sphere. The integral law is also valid for an annular region centered at the origin with both the boundaries within the source or one of the boundary inside the source. The field within the annular region is given by the multipole coefficients evaluated at one of the boundary two-spheres. In eqn.(22) the multipole coefficients depend parametrically on the field coordinates and the application of the integral version of Gauss's divergence law is not a straight-forward issue. For points outside the source Gauss's integral theorem is valid for any annular region surrounding the source.

Lastly, if we consider $\vec{\nabla}$ in terms of the Cartezian coordinates $\vec{\nabla}_{\vec{r}} \equiv \vec{\nabla}_{\vec{R}}$ and this is in general not valid in the spherical polar coordinates while taking the divergence of a vector field. In the Cartezian coordinates $\vec{\nabla}_{\vec{r}} \cdot \vec{E} = \vec{\nabla}_{\vec{R}} \cdot \vec{E}$ and hence $\vec{\nabla}_{\vec{r}} \cdot \vec{E} = \frac{q}{\epsilon_0} \delta(\vec{r} - \vec{r}')$ for the inverse square field of a point charge not at the origin. This is not valid, as is evident from eqns (73, 78), if we take the divergence using the spherical polar coordinates and these aspects are not considered in standard literature, e.g, equations: (6.58, 6.59) [23]. In fact the the inverse square law is not the solution of the following equation expressed in terms of (r, θ, ϕ) :

$$\vec{\nabla}_{\vec{r}} \cdot \vec{E} = \frac{q\delta(\vec{r} - \vec{a})}{\epsilon_0} \quad (85)$$

and the solution of the above equation, if exists, is not physically significant. As we have discussed earlier only for a point charge at the origin the above Green's function equation has a solution given by $V(r) = \frac{q}{4\pi\epsilon_0 r}$. For a point charge not at the origin the potential is given in terms of the Legendre polynomials and satisfies the homogeneous Laplace's equation which is a local differential equation. As we have mentioned below eqn.(71) the potential is well-defined for all points apart from $\vec{r} = \pm\vec{r}'$. The potential given by the Legendre polynomials is singular at $\vec{r} = \vec{r}'$. The potential at $\vec{r} = -\vec{r}'$ is not well-defined. However both the above problems are not physically significant in a quantum mechanical description of charge densities or if we take into account the finite volume of the elementary particles so that the charge densities are finite and satisfy the regularity conditions discussed below eqn.(71). For non-singular sources the force law in the spherical polar coordinates (r, θ, ϕ) is given by eqn.(84) and for a non-singular source this expression is well-behaved for all points. Equation (83) is not the solution of eqn.(85) and satisfies the homogeneous Laplace's equation although the total flux of the electric field is non-vanishing for any sphere with $r > r'$. This is not contradictory to the integral version of the Gauss's law. We should note that the electric field is singular at the point charge and as we have mentioned earlier, in general, it is not possible to apply Gauss's integral law to singular fields. We can apply Gauss's integral law to the corresponding electric field for any region leaving out the point charge. In the spherical polar coordinates (r, θ, ϕ) we have an integral version of the Gauss's divergence theorem for a point charge only when the point charge is at the origin. An analogous statement similar to the integral law is also valid for the higher order pure multipole fields like the dipole field although the fields are singular at the origin. This is due to the fact that the pure multipole fields are

regular everywhere apart from the origin and the integral law is valid for any annular region surrounding the origin. The divergences of the pure multipole fields are vanishing everywhere. The fields are regular everywhere apart from the origin and since the radial dependence is of the form $1/r^2$; $n \geq 3$ the angular dependence are such that the integral theorem remains to be valid for any annular region centered at the origin.

We conclude our discussions on Maxwell's equations for steady sources with a few discussions on the derivation of the curl of $\vec{B}$ law. In the derivation of $\vec{\nabla} \times \vec{B}$ law we have a 'boundary' term [24]:

$$I_B = \int \vec{\nabla}_{\vec{r}'} \cdot \left[\frac{\vec{J}(\vec{r}')}{R} \right] dv' \quad (86)$$

We can not use Gauss's divergence theorem directly as $\frac{\vec{J}(\vec{r}')}{R}$ is divergent at $\vec{R} = 0$. We break the integral into two parts:

$$I_B = \int_{\delta v} \vec{\nabla}_{\vec{r}'} \cdot \left[\frac{\vec{J}(\vec{r}')}{R} \right] dv' + \int_{V-\delta v} \vec{\nabla}_{\vec{r}'} \cdot \left[\frac{\vec{J}(\vec{r}')}{R} \right] dv'$$

Here δv is an arbitrarily small sphere surrounding $\vec{r}$. For a regular current distribution the first term vanishes in the limit $R \rightarrow 0$. In the second term $\frac{\vec{J}(\vec{r}')}{R}$ is well-behaved every where within $V - \delta v$. We can use the divergence theorem to transform the volume integral into two boundary surface integrals. For the inner boundary $\int \frac{\vec{J}(\vec{r}')}{R} \cdot d\vec{s}_1$ is vanishing for a regular current distribution as the radius of δv become arbitrarily small. For the outer boundary $\vec{J} \cdot d\vec{s}_2$ is zero as the current should be tangential at the boundary of the source. However this derivation is valid when the derivatives of $\vec{J}$ are well-behaved within the source apart from the boundary. Similar discussions are valid in the spherical polar coordinates with $(\frac{1}{R})$ is replaced by the expansion in terms of the Legendre polynomials. As we have mentioned earlier we can interchange the variables $\vec{r}$ and $\vec{r}'$ at the cost of a minus sign while taking the gradient of the potential (83) and together with the required regularities of the source similar to those mentioned above we have $\vec{\nabla}_{\vec{r}} \cdot \vec{A} = 0$ in the spherical polar coordinates. We should note that the above discussions are valid only for steady currents with the required regularities.

We can prove the divergence of $\vec{B}$ law from the fact that the magnetic field is the curl of the vector potential as is obvious from the Biot-Savart's law. However in the spherical polar coordinates the magnetic vector potential should be expressed in terms of the Legendre polynomials. It can be shown from the multipole expansion of the magnetic vector potential that there exists no magnetic monopole field [24].

We now consider the no-work law for the electric field. For regular sources the arguments will be similar as above for the magnetic field so that we can apply Stoke's law to prove the no-work law for the electric field. For a point charge we can consider an annular region surrounding the point charge with the inner boundary being an infinitesimally small circle centered at the point charge. The work done for this inner boundary vanishes out of spherical symmetry and we have the no work law for the outer closed line. In the spherical polar coordinates we can consider an additional infinitesimally small circle centered at the south-pole w.r.t point charge. The work done for this circle vanishes in the limit the radius of this circle vanishes and we again have the no-work law for the electric field.

In the above discussions in this section on electrodynamics we have considered the potentials and the fields with static sources. We have found that with the inverse square laws the electric and the magnetic potentials do not obey the corresponding Poisson's equations in terms of the spherical polar coordinates. In terms of the spherical polar coordinates the Laplacian of $(\frac{1}{R})$ do not contain a delta function and give non-local terms. Following experiments we have assumed that the potentials for a point charge not at the origin is given in terms of the Legendre polynomials, equation (83). Equation (83) satisfy the homogeneous Laplace's equation and $(\frac{1}{R})$ is not the generating function for the expression (83) for all points. Similar discussions are valid for the magnetic vector potentials for steady currents satisfying the regularity conditions mentioned below equation (86). In the static situation of static charge density and static steady currents we have the following equations for the potential and the fields in the spherical polar coordinates:

$$\nabla^2 V = 0; \quad \nabla^2 \vec{A} = 0$$

the fields are given by: $\vec{E} = -\vec{\nabla}V$ and $\vec{B} = \vec{\nabla} \times \vec{A}$. In the spherical polar coordinates all the Maxwell's equations remain valid only in the homogeneous form. When time varying steady currents are present we will have to consider Faraday's law. The potential formulation will be consistent with the Faraday's law if we redefine the fields in terms of the potentials in the following way: $\vec{E} = -\vec{\nabla}V - \partial_t \vec{A}$ and $\vec{B} = \vec{\nabla} \times \vec{A}$. However Maxwell's first and fourth equations, even in the homogeneous form $\vec{\nabla} \cdot \vec{E} = 0$ and $\vec{\nabla} \times \vec{B} = \mu_0 \epsilon_0 \partial_t \vec{E}$ may not remain valid in the spherical polar coordinates for time varying steady currents which give electromagnetic radiation. We discuss this situation in the following paragraph.

We now consider the most general situation of non-steady currents. To discuss the Maxwell equations for the potentials and the fields we will consider the radiation from an oscillating point charge in terms of the spherical polar coordinates in Supplement:X. We first discuss the conventional treatment in terms of a claimed Green's function of the wave equation in the spherical polar coordinates. We will discuss in Supplement:X that the Green's function, eqn (120), is not the solution of the corresponding inhomogeneous wave equation. This is similar to the discussions in the static situations below eqn (79). The resulting expression for the potential and the fields do not obey the wave equations. Also the Lorentz gauge condition is violated. Thus the corresponding potentials and the fields are not the conjugate pair of harmonics in the spherical polar coordinates. We resolve this problem with the help of a second quantization scheme and the fact that photons travel in null geodesics in vacuum. We assume that there exist a scalar electric potential and a magnetic vector potential obeying the homogeneous wave equations in the spherical polar coordinates. In a second quantization scheme the modal solutions of the above wave equations creates photons traveling along the null geodesics in vacuum. The potential for a general source is obtained from the distribution formed in terms of the modal solutions of the wave equations for the potential. This is similar to the corresponding static situation where the potentials in the spherical polar coordinates are given by the distributions formed out of the normal modes of the homogeneous Laplace's equation expressed in terms of the Legendre polynomials. We will give a few modal solutions of the wave equation in the spherical polar coordinates. The electromagnetic fields are related to the potentials by the relations similar to the above paragraph: $\vec{E} = -\vec{\nabla}V - \partial_t \vec{A}$ and $\vec{B} = \vec{\nabla} \times \vec{A}$. If the fields thus obtained would have satisfied the wave equations we would have a conjugate pair of harmonics in the spherical polar coordinates and the Lorentz gauge would also have been satisfied. In the spherical polar coordinates in four dimensions this appears not to be the situation. The resulting fields do not obey the wave equations although in terms of a more fundamental description, namely the corresponding quantum field theory, nothing acausal happens. The photons created by the modal potentials travel with the velocity of light. The well-known Maxwell's first and fourth equations for non-steady currents are not valid in the spherical polar coordinates. The most appealing aspect of the above formulation for time dependent sources is that the equations for the potentials reduce to the corresponding static equations in the static limit as we have discussed in Supplement:X.

Thus in flat space-time we can not obtain the potentials and the fields in terms of the spherical polar coordinates by changing the coordinates in the expressions for the potentials and the fields expressed in terms of the Cartesian coordinates. We should solve the equations for the potentials expressed in terms of the spherical polar coordinates to obtain the potentials and the fields in terms of the spherical polar coordinates. For the time independent sources we do not have the corresponding inhomogeneous equations in terms of the spherical polar coordinates. We write the corresponding homogeneous equations in terms of the spherical polar coordinates. We thereafter solve the homogeneous equations for both points inside and outside the source. This remains to be valid for source densities not involving the fields themselves unlike sources following the Ohm's law. For points inside the source the permittivity and the permeability will be different from that of the vacuum. The relationship between the source and the solution is obtained from the globally defined multipole coefficients. This is not similar to the corresponding situation in the Cartesian coordinates where the fields are given by the solutions of the corresponding inhomogeneous equations with the locally defined charge-current densities as sources. For time dependent steady currents or for non-steady currents we may not have Maxwell's first and fourth equation and adopt the potentials satisfying the homogeneous wave equations as fundamental physical variables. The fields are obtained from the potentials through the following relations: $\vec{E} = -\vec{\nabla}V - \partial_t \vec{A}$ and $\vec{B} = \vec{\nabla} \times \vec{A}$.

The above discussions may also be significant for field theory in curved space-time where the scale factors within the divergence operator and the partial derivatives of R w.r.t $\vec{r}$ and $\vec{r}'$ may not have any definite symmetry. In these situations we follow the above procedures to describe the electromagnetic fields in terms of the potentials. Even if the electromagnetic field in a particular curved space-time is given by the inhomogeneous equations the discussions in this article demonstrate that the diffeomorphism invariance of the inhomogeneous equations is a non-trivial issue and photons travel along null geodesics in vacuum is a more fundamental principle in this respect. In passing we note that the electric potential outside the source of a spherically symmetric asymptotically flat charged static space time is given by the equation: $\frac{\partial V}{\partial r} = \frac{Qh_r}{r^2}$. Here ' Q ' is the charge as is measured from the field at infinity and $g_{rr} = h_r^2$.

The above discussions are also significant for constructing the Lagrangian of the electromagnetic fields for both curvilinear coordinates and in curved space-time. In particular the Lagrangian of the electromagnetic field in the spherical polar coordinates in the static situation is given by only the free field homogeneous part. In the dynamical situation the Lagrangian density in the spherical polar coordinates is expressed in terms of the four potentials: $A_\mu = (\frac{V}{c}, \vec{A})$ and is given by the expression: $-\frac{c}{\mu_0} A_{\alpha,\beta} A^{\alpha,\beta}$. The stress tensor is expressed in terms of the four potentials and is to be obtained from the Lagrangian. This is significant

in the context of the corresponding quantum field theory. Although the lagrangian does not contain the interaction term explicitly, in a second quantization scheme, the solutions in the dynamical situations [eqns (127..)] are similar to the interaction action in the Cartezian coordinates and gives the coupling between the source and the electromagnetic field. Only in the special situation when the point charge is at the origin the expression $\frac{q}{r^2}\hat{r}$ remains to be the solution of the corresponding Poisson's equation expressed in terms of the spherical polar coordinates and the Lagrangian will contain the source density.

We conclude this section with the following discussions on the electrostatic behavior of conductors. We should note that with an ideal surface charge density the electric field within an arbitrarily-shaped conductor may not vanish. This follows from Coulomb's law. We can consider a simple situation of a cube with sides of length $2a$ and is centered at the origin. Firstly, the externally given charge should be distributed equally on the six surfaces and the surface charge densities on the surfaces should be the same function w.r.t their respective centers. Secondly, the charge distribution on the surfaces should be a symmetric function under reflections w.r.t the centers of the respective surfaces. The surface configuration is symmetric under reflection w.r.t the coordinate planes and the field together with the surface charge densities on the top or the bottom surfaces can not change if we just rotate the cube about the Z -axis by an angle π . Let us now consider a point on the Z -axis with coordinates $(0, 0, \epsilon)$ where $\frac{\epsilon}{a} \rightarrow 0$. If the field has to vanish at ' P ' we will have the following condition on the surface charge density $\sigma(u, v)$: $\int \int f(u, v) du dv = \frac{1}{a^2} \int \int (u^2 + v^2) f(u, v) du dv$, where $f(u, v) = \frac{\sigma(u, v)}{(a^2 + u^2 + v^2)^{\frac{3}{2}}}$ and (u, v) are the orthogonal Cartesian coordinates on the surfaces w.r.t their centers. One can obtain the $(u^2 + v^2)$ factor in the second integral by considering rotations about the X - and Y - axes by angle $\frac{\pi}{2}$ together with the symmetries of the charge distributions on the surfaces. Clearly it is not possible to find a non-trivial surface charge density satisfying the above condition.

We now give a more realistic explanation of the electrostatic behavior of a charged conductor. We consider an arbitrarily-shaped thin metallic shell. If we do not represent the charges at the outside surface by an ideal surface charge density (which is the situation in reality) there will be higher order multipole fields which are directed inwards for electrons. To the leading order the multipole potential is of the form $r \int \rho(\vec{r}') \cos(\gamma) dr' d\Omega'$. The exact form of the multipole potential will depend on the nature and the symmetry of the source. The corresponding multipole force will push the free electrons within the core towards the inner surface. However this will create a small positively charged core in between. This core will impart an attractive force on the electrons and ultimately an equilibrium condition will be reached. If the multipole force is attractive then also a layer of positive charges will be created near the surface as the electrons in this layer are attracted towards the boundary and again an equilibrium will be reached. The field within the region enclosed by the shell will be almost vanishing apart from microscopic fluctuations near the outer surface of the shell. This whole distribution of the charges can take place within a few atom thick layer at the outer surface of the shell. These discussions indicate that any small departure from the inverse-square nature may not be observed in experiments similar to those discussed in the introduction of [23]. Thus the electric field can have a term of the form $\alpha(r) \frac{\hat{r}}{r^n}$ where $\alpha(r) \ll 1$ and $n > 2$ with the force within a conductor vanishing. In this case it is not possible, in general, to have a Poisson's equation. Also these correction terms will introduce non-locality. For points outside the sources we will have an equation different from the Laplace's equation. However these terms may be significant to explain charge screening.

We should also note that Legendre polynomials are either symmetric or antisymmetric around $\theta = \frac{\pi}{2}$ and for a charge distribution on an arbitrary shaped conductor which can not be expressed as a sum of a symmetric and an antisymmetric part we can not expand the corresponding potential in terms of Legendre polynomials.

We can construct a well behaved current density for the elementary charge carriers: $\vec{J} = \rho([r - r_c(t)]) \vec{v}_c(t)$. Here we have considered the charge density of the elementary charge carriers to be spherically symmetric and well-behaved, vanishing smoothly to zero around the boundary. The particle is a sphere of radius a and we can have a local charge conservation law leading to Maxwell's equations.

Faraday's law, with the following convention (Section:7.1.3 [24]),

for motional emf or induced electric field the direction of the current or the electric field along a closed loop and the orientation of the enclosed surface giving the magnetic flux are related by the right-hand thumb rule,

together with the above discussions and the current density equation (differential version of the electric charge conservation law) reproduces Maxwell's laws of Classical Electrodynamics.

We conclude our discussions on the Maxwell's equations with a few comments on the forms of the Maxwell's equations in the spherical polar coordinates, (r, θ, ϕ) , with sources depending explicitly on the fields. We consider a source obeying the Ohm's law, $\vec{J} = \sigma \vec{E}$. In this case $\vec{\nabla} \cdot \vec{E}$ has to be non-vanishing out of the charge conservation law and is given by: $\vec{\nabla} \cdot \vec{E} = -\frac{1}{\sigma} \frac{\partial \rho}{\partial t}$. In this case the fourth equation in the spherical polar

coordinates, $\vec{\nabla} \times \vec{B} = \mu\epsilon \frac{\partial \vec{E}}{\partial t}$ leads to contradiction if we take the divergence of both sides. Since $\vec{J}$ depends explicitly on the field $\vec{E}$ we follow the corresponding inhomogeneous equation and the fourth equation, in terms of the spherical polar coordinates, is given by:

$$\vec{\nabla} \times \vec{B} = \mu\sigma \vec{E} + \mu\epsilon \frac{\partial \vec{E}}{\partial t}$$

This leads to the following expression for the charge density if we take the divergence of both sides:

$$\rho(t) = \rho(0)\exp(-\frac{\sigma}{\epsilon}t)$$

and the first equation becomes,

$$\vec{\nabla} \cdot \vec{E} = \frac{1}{\epsilon}\rho(0)\exp(-\frac{\sigma}{\epsilon}t)$$

in accordance with the corresponding equation in the Cartesian coordinates [24].

We now make some comments on ideal surface and line current densities. Let us first consider a surface with an edge, i.e., two surfaces joined along a curve making an angle which can, in general, vary along the curve. If a surface current density originates at one surface the current can not propagate to the second surface as the only way that the current can have vanishing perpendicular components to both the surfaces is to propagate along the edge (geometrically for two surfaces making an edge the only way that the ideal surface current density can remain tangential to both the surfaces at the edge is to flow along the edge). The above arguments demonstrates that we can not have surface currents out of point charges flowing along an arbitrarily shaped surface as an arbitrary surface can be expressed as a collection of infinitesimal flat surfaces with non-parallel normals. Similar arguments as above demonstrates that we can not have currents in an arbitrary curve out of motion of freely flowing point charge carriers unless one applies forces everywhere tangential to the surfaces/curves.

After that we have reestablished Maxwell's equations we should note that the finite volume of the elementary charge carriers indicates that ideal line/surface charge densities cannot exist in nature unless the charge carriers can move with velocity c in one or two directions. Also to have ideal line/surface current densities the charge carriers should move with speed greater than c . Point charges can not exist in nature and the electromagnetic self-energies of the elementary particles are not infinite.

We will now consider the following problem:

We can put a point charge $Q = -\frac{5}{4}q$ at the center of a square with four point charges q at the vertices. The system will be in electrostatic equilibrium. Similar static equilibrium can be defined with three collinear point charges $q, q, -q/4$ with $-q/4$ midway. It can be easily shown that for any other moving observer (moving with velocity V w.r.t the rest frame of the five charges) this system will not be in equilibrium (the electric forces won't balance each other and there will also be magnetic forces) and the point charges will be in relative motion with respect to each other. The magnetic forces being smaller by a factor $\frac{v^2}{c^2}$ can not balance the electric forces. Thus in the moving frame the configuration is not in equilibrium and the charges will either fly away to infinity or collapse making the system non-existent in the moving frame. Similar situation will arise with three charges if one moves in a non-collinear direction. However the above discussions are not significant in the microscopic regime because of quantum mechanical uncertainty principle and charge screening.

We conclude this section with a few comments on magnetic monopoles in the Cartesian coordinate system. It is straight forward to show that under a general electromagnetic duality transformation, eqn.(6.151, 6.152) [23],

$$\vec{E} = \vec{E}'\cos\xi + \vec{H}'\sin\xi \quad \vec{D} = \vec{D}'\cos\xi + \vec{B}'\sin\xi$$

$$\vec{H} = -\vec{E}'\sin\xi + \vec{H}'\cos\xi \quad \vec{B} = -\vec{D}'\sin\xi + \vec{B}'\cos\xi$$

and

$$\rho_e = \rho'_e\cos\xi + \rho'_m\sin\xi \quad \vec{J}_e = \vec{J}'_e\cos\xi + \vec{J}'_m\sin\xi$$

$$\rho_m = -\rho'_e\sin\xi + \rho'_m\cos\xi \quad \vec{J}_m = -\vec{J}'_e\sin\xi + \vec{J}'_m\cos\xi$$

the differential version of the magnetic charge conservation law does not remain time-reversal symmetric due to the pseudoscalar and pseudovector nature of magnetic monopole charge and magnetic current density vector respectively, i.e., if $\vec{\nabla} \cdot \vec{J}_m = -\frac{\partial \rho_m}{\partial t}$ is time-reversal symmetric then $\vec{\nabla} \cdot \vec{J}_m = -\frac{\partial \rho_m}{\partial t}$ no-longer remains time-reversal symmetric although the differential version of the electric charge conservation law remains time-reversal symmetric under electromagnetic duality transformation.

XIV. SUPPLEMENTIV: COMMENTS ON HYDRODYNAMICS

In this article we will review the laws of fluid dynamics. Our discussions will be based on mainly that of chapter 40, 41 of The Feynman Lectures on Physics, Vol.2 [25].

The dynamics of dry water is governed by eqn.(40.6) [25]:

$$\frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla}) \vec{v} = -\frac{\vec{\nabla} p}{\rho} - \vec{\nabla} \phi \quad (87)$$

or using a vector analysis identity to the second term of the above equation:

$$\frac{\partial \vec{v}}{\partial t} + (\vec{\nabla} \times \vec{v}) \times \vec{v} = -\frac{\vec{\nabla} p}{\rho} - \vec{\nabla} \phi \quad (88)$$

where $\vec{v}$ is the velocity of an fluid element for which such laws can be applicable, p is the fluid pressure and ϕ is the potential per unit mass for any potential force present. We can derive some important laws from eqn.(86). The first one is the equation for vorticity ($\vec{\Omega} = \vec{\nabla} \times \vec{v}$) and is obtained by taking curl of eqn.(87):

$$\frac{\partial \vec{\Omega}}{\partial t} + \vec{\nabla} \times (\vec{\Omega} \times \vec{v}) = 0 \quad (89)$$

The second one is Bernoulli's theorems (40.12) and (40.14) [26]:

$$\vec{v} \cdot \vec{\nabla} \left(\frac{p}{\rho} + \phi + \frac{1}{2} v^2 \right) = 0 \quad (90)$$

i.e,

$$\frac{p}{\rho} + \phi + \frac{1}{2} v^2 = \text{const (along streamlines)} \quad (91)$$

valid for steady flow and

$$\frac{p}{\rho} + \phi + \frac{1}{2} v^2 = \text{const (everywhere)} \quad (92)$$

valid for steady and irrotational flow.

However in all these equations the variation of the fluid density, ρ , is not considered while in deriving eqn.(40.17) [25] the variation of fluid density is not properly taken into account. The consideration for variation of the density of a nearly-incompressible fluid may become important through the facts that when unconstrained the shape of a fluid can be changed almost freely and sparsed away and through the facts that layers of fluids can be very easily spread or detached away although these properties vary from fluid to fluid. These features together with the local version of the conservation of mass law (assuming that there is no local source or sink in the region of interest):

$$\vec{\nabla} \cdot (\rho \vec{v}) = 0 \quad (93)$$

indicate that we should consider the possibility for variation of ρ properly as we will illustrate later that some ideal models can cause a finite variation of ρ and in reality the description of the motion should be changed. While the divergence of $\vec{v}$ may become important in cases like Couette flow where the centrifugal forces imposes a finite and may even be large divergence of $\vec{v}$.

We can derive a proper version of eqn.(86) by applying Newtons second law to the fluid momentum per unit volume and we have:

$$\frac{\partial(\rho \vec{v})}{\partial t} + [\vec{v} \cdot \vec{\nabla}](\rho \vec{v}) = -\vec{\nabla} p - \vec{\nabla}(\rho \phi) \quad (94)$$

This equation is in general a non-linear coupled [through eqn.(91)] partial differential equation for $\vec{v}$. Bernoulli's theorems for fluid dynamics can only be established when ρ is constant :

$$\vec{v} \cdot \vec{\nabla} [p + (\phi\rho) + \frac{1}{2}(\rho v^2)] = 0 \quad (95)$$

i.e.,

$$p + (\phi\rho) + \frac{1}{2}(\rho v^2) = \text{const (along streamlines)} \quad (96)$$

valid for steady flow and

$$p + (\phi\rho) + \frac{1}{2}(\rho v^2) = \text{const (everywhere)} \quad (97)$$

valid for steady and irrotational flow.

In general, when ρ is varying, only the first of the Bernoulli's theorems :

$$\vec{v} \cdot \vec{\nabla} [p + (\phi\rho) + \frac{1}{2}(\rho v^2)] = 0 \quad (98)$$

remains to be valid provided $(\vec{v} \cdot \vec{\nabla} \rho)$ is vanishing or is approximately valid if $|(\vec{v} \cdot \vec{\nabla} \rho)\vec{v}|$ is negligible compared to the other terms in eqn.(93). To illustrate the significance of these comments, let us consider the ideal model to calculate the efflux-coefficient, fig. 40-7 [25]. After that the contraction of the cross-section of the emerging jet has stopped we have, from the conservation of mass law, the following equation for ρv at two vertical points:

$$\rho_1 v_1 = \rho_2 v_2 \quad (99)$$

In this case pressure is the atmospheric pressure and remains the same throughout the flow and thus even for the flow of a nearly-incompressible fluid ρ can vary as v changes with height. In reality the flow usually gets sparsely away after a distance which varies for different flows.

The viscous flow of a fluid is governed by the following two laws which are obtained from eqn.(93) and eqn.(41.15),[25]:

$$\frac{\partial(\rho\vec{v})}{\partial t} + [\vec{v} \cdot \vec{\nabla}](\rho\vec{v}) = -\vec{\nabla} p - \vec{\nabla}(\rho\phi) + \eta \nabla^2 \vec{v} + (\eta + \eta') \vec{\nabla}(\vec{\nabla} \cdot \vec{v}) \quad (100)$$

$$\vec{\nabla} \cdot (\rho\vec{v}) = 0 \quad (101)$$

supplemented by proper boundary conditions. To illustrate the significance of the boundary conditions we can consider the change of the shape of the surface of water in a bucket when the bucket is given a steady rotational motion about its axis. The surface of the water become paraboloidal when the bucket is rotating. This shape can not be obtained without a vertical component of fluid velocity along the bucket surface for a finite duration although the bucket surface only have an angular velocity.

In the above equations η is the “first coefficient of viscosity” or the “shear viscosity coefficient” and η' is the “second coefficient of viscosity”. This equation is extremely significant in the sense that this equation, not eqn.(41.16) [26], is the equation which contains all the terms relevant to describe the dynamics of viscous fluids, both nearly-incompressible and compressible. For compressible fluids ρ will also depend on pressure, $p(\vec{r})$. We can modify this equation only through varying the nature of the viscous force.

The equation for vorticity is given by:

$$\begin{aligned} & \frac{\partial \vec{\Omega}}{\partial t} + \vec{\nabla} \times (\vec{\Omega} \times \vec{v}) - \vec{\Omega}(\vec{\nabla} \cdot \vec{v}) + \vec{v} \times \vec{\nabla}(\vec{\nabla} \cdot \vec{v}) = \frac{\eta}{\rho}(\nabla^2 \vec{\Omega}) \\ & - \frac{(\vec{\nabla} p) \times (\vec{\nabla} \rho)}{\rho^2} - \frac{[\vec{\nabla}(\rho\phi)] \times (\vec{\nabla} \rho)}{\rho^2} + \frac{\eta(\nabla^2 \vec{v}) \times (\vec{\nabla} \rho)}{\rho^2} + \frac{\eta + \eta'}{\rho^2} \vec{\nabla}(\vec{\nabla} \cdot \vec{v}) \times (\vec{\nabla} \rho) \end{aligned} \quad (102)$$

We can obtain an equation similar to eqn.(41.17) [25] describing the motion of a viscous fluid past a cylinder provided we can neglect the terms involving $\vec{\nabla}\rho$ and it is given by:

$$\frac{\partial \vec{\Omega}}{\partial t} + \vec{\nabla} \times (\vec{\Omega} \times \vec{v}) - \vec{\Omega}(\vec{\nabla} \cdot \vec{v}) + \vec{v} \times \vec{\nabla}(\vec{\nabla} \cdot \vec{v}) = \frac{\eta}{\rho}(\nabla^2 \vec{\Omega}) \quad (103)$$

Following the procedure in section 41-3,[26] we can rescale the variables to obtain an equation which has Reynolds number (R) as the only free parameter :

$$\frac{\partial \vec{\omega}}{\partial t'} + \vec{\nabla}' \times (\vec{\omega} \times \vec{u}) - \vec{\omega}(\vec{\nabla}' \cdot \vec{u}) + \vec{u} \times \vec{\nabla}'(\vec{\nabla}' \cdot \vec{u}) = \frac{1}{R}(\nabla'^2 \vec{\omega}) \quad (104)$$

where the prime describe the scaled variables, $\vec{u}$ is the scaled velocity and R is given by the usual expression, $R = \frac{uVD}{\nu}$.

To conclude in this section we have derived the exact equation describing fluid dynamics. We considered the motion of both non-viscous and viscous fluids. We proved that in both the cases there are terms which are neglected in the conventional theory but may become significant in some ideal model and in reality the description of motion is changed. Some of these terms even change the dynamical laws of viscous fluid motions by violating the conventional theory established in term of the Reynold number and these terms are significant for the dynamics of compressible fluids like air.

XV. SUPPLEMENTV: A FEW QUESTIONS

What happens to the entropy increase principle as the Universe evolve to form the big-crunch singularity? What happens to the uncertainty relations along the process of gravitational collapses? What is the quantum mechanical description of a Radon atom in a rigid box when the distance of consecutive nodes and antinodes of ψ is equal to or less than the diameter of the atom? What is the position-space wave function of two finite volume massive bosons if we take contact interaction [we can characterize contact interaction in a relativistic covariant way if we take $\Delta x, \Delta t = 0$, i.e, do not consider the coordinates of the center of the particles to characterize the interaction kinematically] into account? How a photon produce electron-positron pair with finite volume concentrate rest masses? What are the charges and masses of the electron-positron pairs forming loops in the vacuum? How two particles with three-momentums k_1, k_2 ($k_1 \neq k_2$) produced to form a loop at a space-time point always arrive at another spacial point simultaneously? What is the microscopic explanation in terms of particle exchanges of the force in the Casimir effect? (If a loop begin at one plate and terminate at the other then the pairs, being tangential, can not impart any energy-momentum to the plates. This is also relevant for the discussions at the beginning of this article and discussions regarding bound states in [38]) What is the mechanism of the collapse of the momentum-space wave function of a particle knocking out an electron from an atom? What is meant by $|\Psi\rangle = c_1(t)|\Psi_{U^{238}}\rangle + c_2(t)|\Psi_{Th^{234}}\rangle$? Is the radiation out of a single atomic transition a spherical wave? Quantum mechanically the region between the rigid walls (which is equiprobable in classical mechanics) is non-homogeneous for a particle in a rigid box ! A single photon can not reproduce Maxwell's equations apart from moving with velocity c . How can a process involving only a few photons be described starting from the Maxwell's equations? The large scale structure of the Universe is homogeneous.

XVI. SUPPLEMENTVI: A FEW COMMENTS ON VIEWING

Let viewing be a relative process which is achieved by comparing the images of the surroundings formed at the brain with the images formed at the brains reduced by a factor N . We can scale time-intervals without measuring any spatial length. Time-intervals are primarily and fundamentally determined by our brains through the axial rotation of the Earth and do not involve any change at our brain similar to the spatial extensions of objects while seeing. Consequently the velocities and accelerations of the images of particles and signals at our brains are reduced by a factor N . Thus the images (and their dynamics) of the surroundings at our brains are not an exact replica of the surroundings. Relating the surroundings with the corresponding images formed at our brain is a part of mind/intelligence.

Is the images formed in our brain are always reduced in size?

One can add to this question through elementary observations on babies. In this context we should also note that forming larger images is essential to a have developed distant vision.

We note that with a sharp resolution (10^{-7}m if we consider that we can see a 0.1mm thick hair at 1m away and the focal length of the eye-lens to 1mm) the processing of image at the retina can involve only one cell.

XVII. SUPPLEMENTVII: A FEW COMMENTS ON UNCERTAINTY PRINCIPLE AND RADIATIONS INSIDE ISOTHERMAL ENCLOSURES

In this section we first discuss the position-momentum uncertainty relation in the context of optical microscope observations. For microscope observations if we analyse the scattering of a photon by an electron through Compton scattering then to the leading order the uncertainty in the x -component of the electron momentum for photon wavelengths greater than twice the Compton wave-length of the electron is

$$\Delta p_x = \frac{2h \cos \phi}{\lambda} - \frac{2h^2}{m_0 c} \frac{1}{\lambda^2} \cos \phi \quad (105)$$

Here ϕ is the angle of scattering of the photon in the forward direction and the photon is assumed to be incident along the positive X -direction ($\pi/2 - \phi$ is the angle of incidence of the scattered photon w.r.t the microscope lence). The maximum uncertainty in p_x of the electron, in the context of Compton scattering, that the electron is observed happens for $\phi = 0$.

Consequently the approximate minimum position-momentum uncertainty relation becomes:

$$\Delta x \Delta p_x = h - \frac{h^2}{m_0 c \lambda} \quad (106)$$

The expression (for $\phi = 0$) valid for all λ is:

$$\Delta x \Delta p_x = \frac{1.22h}{n} - \frac{1.22h}{2n} \frac{2\lambda_C}{\lambda + 2\lambda_C} \quad (107)$$

Here λ_C is the Compton wave-length of the electron.

The r.h.s varies with λ and for $\lambda \gg \lambda_C$ the right hand side of the minimum uncertainty product is $\frac{1.22h}{n}$. The above equation gives the minimum uncertainty product in the sense that we have considered the maximum resolution for the incident photon with wave-length λ and this leads, in principle, to a value $\frac{1.22h}{2n}$ for the r.h.s when $\lambda \ll \lambda_C$.

In general the right-hand side also depends on $\cos \phi$.

We now discuss a few aspects of radiations inside isothermal enclosures. According to Wien's law [30] the radiation inside a hollow iron isothermal enclosure at it's red-heat temperature is peaked about a wave-length which is about five times larger than that of visible light and the radiation in the visible spectrum may be negligible.

Are the radiations inside a slate-isothermal enclosure and an iron-isothermal enclosure the same at the red-heat temperature of iron.

In the derivation of Rayley-Jeans law for radiation inside a cubic isothermal enclosure we have the following relation between the allowed set of modes (with perfectly reflecting boundary conditions) for two cubic enclosure of sides L' and L respectively:

$$\frac{L'}{L} = \sqrt{\frac{n'_x{}^2 + n'_y{}^2 + n'_z{}^2}{n_x{}^2 + n_y{}^2 + n_z{}^2}} \quad (108)$$

Now if there exists a radiation within the second cube for a particular set of values of (n_x, n_y, n_z) the above equation in general may not give integral set of solutions for (n'_x, n'_y, n'_z) for arbitrary values of L and L' . Thus the corresponding mode of radiation will be absent in the first cube, something which is unexpected from the radiation mechanism through atomic structure with the material remaining the same.

The above discussions are also relevant for the equilibrium quantum statistical mechanical description, in the context of varying the volume of the enclosure, of particles in a rigid box.

We now consider an interesting paradox. We can consider two absorbers inside an isothermal enclosure. The absorbers can only absorb radiation with two different bands of wave-lengths and pass the rest. Since e_λ is different we can construct a Carnot-engine similar to those discussed in page-118 [30] violating the second law of thermodynamics. However in reality this do not lead to any contradiction. Let us consider the process in detail. Let the absorbers be parallel to two faces of the cube. Due to the presence of the absorbers there will be tubes of radiation with a given range of wave-length (the portions that the absorbers absorb) that wont be reaching corresponding area-elements of the parallel surfaces. But these area-elements will keep on radiating the corresponding radiations. This will remain valid at any stage. Thus the temperature of the enclosure will keep on decreasing and it is not possible to operate the Carnot-engine without changing anything else and thereby violate the second law of thermodynamics. These arguments are also applicable for the Carnot-engines considered in page-118 [30]. In this context he perfectly reflecting mirror models are misleading.

We will later discuss a few aspects of the quantum statistical concept of entropy for an ideal gas kept inside an isolated enclosure taking the collision interactions into account.

In passing we note that the expression $Ne^{i(\omega t - kx)}$ leading to negative energy do not satisfy the non-relativistic Schrodinger equation. We will discuss later the relevance of this aspect in the context of the convergence of the perturbation coefficients with time for perturbing potentials sinusoidal in time.

XVIII. SUPPLEMENT VIII: A FEW COMMENTS ON NEWTONIAN MECHANICS

In this section we will make a few comments regarding Newton's laws of motion for conservative force field.

We define a conservative force to be a force for which the infinitesimal work done by the force field is total derivative, i.e,

$$\vec{F} \cdot d\vec{r} = -dV \quad (109)$$

Here $d\vec{r}$ is an infinitesimal element on the trajectory of the particle in the given force field and is determined by the definition of force in terms of the velocity and its time-derivatives.

The l.h.s of the above equation, $\vec{F} \cdot d\vec{r}$, will be a total derivative if $\vec{F}$ become either $m \frac{d}{dt}(\vec{v})$ or $\alpha_n [\frac{d}{dt}(\vec{v} \cdot \vec{v})^n](\vec{v} \cdot \vec{v})^{n-1} \vec{v}$.

The first choice gives us Newton's second law.

Let us consider the terms of the second category. These terms do not take into account the vectorial nature of velocity completely. The force remains to be vanishing either if the velocity is zero or if the magnitude of the velocity remains constant. If the velocity only changes its direction then this definition of force leads us to conclude that there is no force acting on the particle. The simplest example is the rotational motion. Thus these definitions are not in accordance with the first law of motion.

Another possible class of total integrals are of the form $\beta \frac{d}{dt} [\frac{d\vec{v}}{dt} \cdot \vec{v}] \frac{d\vec{v}}{dt}$. Again the terms inside the parentheses are of the form $\frac{d}{dt}(\vec{v} \cdot \vec{v})$ and the force will again be vanishing if only the magnitude of the velocity remains constant.

We wont consider terms involving time-derivative of the acceleration. The force will then be vanishing for constant accelerations. Also we wont consider $\vec{r}$ dependent terms as these terms will identically lead to zero force either at the origin or at infinity even if the acceleration is non-vanishing at the corresponding points.

If one can show that the definitions of force as are considered in this section are the only possibilities so that $\vec{F} \cdot d\vec{r}$ is a total derivative and the first law remains valid, then we can say that the second law follows from the first law at least for conservative force fields. We should also note we can not define force to be a sum of two terms one of which satisfies the above criteria but the other does not.

In passing we should note that the equation $\vec{F} = -\vec{\nabla}V(x, y, z)$ is valid only for the solutions of the above equation joining two points in a given interval of time. Other trajectories give us different accelerations and thereby different forces acting on the particle. If we do not define force to be mass times the acceleration the definition of kinetic energy will be different. We will later consider the significance of these discussions in the context of the Lagrangian formulation of classical mechanics.

We should note that piecewise discontinuous curves involves impulsive forces and one should take into account this aspect while discussing conservative forces.

XIX. SUPPLEMENT IX: A FEW COMMENTS ON THE ENTROPY OF A SCALAR FIELD IN SCHWARZSCHILD BLACK HOLE BACKGROUND AND THE BRICK WALL MODEL

The thermodynamical nature of black holes was first established by Bekenstein [39]. He obtained an expression for the entropy of black holes. This expression was proportional to the area of the horizons of the black holes. Hawking obtained an exact expression for the entropy by considering the behavior of matter fields in the black hole backgrounds [40]. The entropy of a black hole, considered as a thermodynamical system, was found out to be $\frac{A}{4\pi}$. Here A is the horizon surface area. The thermodynamical properties of matter fields in black hole backgrounds was discussed by 't Hooft [29]. The black hole is assumed to be in thermal equilibrium with surrounding matter. For a static black hole the rate of particles radiated by the black hole should be equal to the rate of absorption of matter by the black hole. 't Hooft [29] assumed that the matter field wave function is vanishing near the horizon as well as at infinity, i.e, for a large value of the radial coordinate. Since then a lot of works had been done on the entropy of matter fields in black hole backgrounds using the brick-wall model [41,42,43,44].

In this section we will discuss a few aspects on the entropy of a massless scalar field in the Schwarzschild black hole background using the brick-wall model of 't Hooft [29]. We will show that with a proper counting of the density of states, the entropy of a massless scalar field is different than that obtained by 't Hooft.

The radial part of the wave equation of a massive scalar field in the Schwarzschild black hole background is given by the following expression:

$$(1 - \frac{2M}{r})E^2\psi(r) + \frac{1}{r^2}\partial_r r(r - 2M)\partial_r \psi(r) - [\frac{l(l+1)}{r^2} + m^2]\psi(r) = 0 \quad (110)$$

Here l is the angular momentum quantum number and E is the energy.

We follow the procedure of 't Hooft to calculate the entropy of a scalar field using the brick wall model and the WKB quantization rule [29]. We consider massless fields but the discussions can also be extended for massive fields. The boundary conditions on the scalar field are the following:

$$\psi(2M + h), \psi(L) = 0 \quad (111)$$

where $h \ll 2M$ and $L \gg 2M$. Here h is the brick-wall cut-off parameter. We consider the solutions of the wave-equation obtained by the WKB approximation. The solutions are assumed to be of the form $\rho(r)\exp iS(r)$. The solutions are stationary w.r.t the radial variable throughout the spatial manifold. These solutions are also important for the analysis of the Hawking radiation.

The expression for the free energy of a massless scalar field is given by the following expression:

$$\pi\beta F = -\beta \int \frac{dE}{[\exp(\beta E) - 1]} \int \frac{dr}{V(r)} \int dl(2l+1) \sqrt{E^2 - \frac{V(r)}{r^2}} [l(l+1)] \quad (112)$$

Here $V(r) = 1 - \frac{2M}{r}$. The range of E integration is from zero to infinity although one should be considering the back reaction to the metric. The range of r integration is from $2M + h$ to L where $L \gg 2M$.

We now consider the free-energy calculation using a proper density of states. Let us first consider the WKB quantization condition together with the brick wall boundary condition. We have the following condition:

$$\pi n = \int_{r_1}^L dr k(r, l, E) \quad (113)$$

Here the symbols have their usual meaning. We assume that for a given set of values for E , l , r_1 and L there exist one solution of the above integral equation. The upper value of l is restricted by the condition that $k(r)$ should remain real throughout the total region of integration. We have,

$$k(r) = \frac{1}{V(r)} \sqrt{E^2 - \frac{V(r)}{r^2}} l(l+1) \quad (114)$$

Here we are considering a massless scalar field. At $r = 2M + h$ with $h \rightarrow 0$, l can be varied freely. However for a state for which $k(r, E, l)$ is real throughout the complete region from $r = 2M + h$ to $r = L$ the upper

limit of l is restricted by the maximum value of $\frac{V(r)}{r^2}$ which occurs at $r = 3M$. Hence the upper limit of l is given by the following expression,

$$l(l+1) = 27E^2M^2 \quad (115)$$

The l integration in eqn.(110) for the free energy introduces a factor $\frac{r^2}{V(r)}$ in the radial integral. We can take the upper limit of l -integration to be $l(l+1) = s(27E^2M^2)$, where s can be taken to be arbitrarily close to one but not one so that $k(r)$ is real for all values of r . The product $s\frac{V(r)}{r^2}(27E^2M^2)$ is less than E^2 for all values of r . We first do the l -integration. The free energy is given by the following expression:

$$\pi\beta F = -\frac{2}{3}\beta \int \frac{dE}{[\exp(\beta E) - 1]} \int \frac{r^2 dr}{V^2(r)} [E^3 - \{E^2 - s(27E^2M^2)\frac{V(r)}{r^2}\}^{\frac{3}{2}}] \quad (116)$$

We can do a binomial expansion in s to find out the leading divergent term in the brick-wall cut-off parameter. The maximum value of the term $27M^2\frac{V(r)}{r^2}$ is one. We thereafter do the radial integration and the energy integration respectively. It can easily be seen the term associated with the linear order term in s is the most divergent term in the brick-wall cut-off parameter h . The entropy is given by $S = \beta^2 \frac{dF}{d\beta}$. The entropy associated with the linear order term in ' s ' is given by,

$$S = \frac{27sI}{128\pi^4} \left[\frac{L}{M} + 2\ln\left(\frac{L}{h}\right) \right] \quad (117)$$

Where $I = \int_0^\infty \frac{x^3 dx}{(\exp x - 1)} = \frac{\pi^4}{15}$. The divergent part of the entropy associated with the brick-wall cut-off parameter is logarithmically divergent in the cut-off parameter. This is expected as the constant-time foliations intersect at the horizon. Here β is given by the usual expression $\beta = 8\pi M$ and h is the brick-wall cut-off parameter. We have assumed $L \gg 2M$. We find that with this expression of β the divergent part of the entropy in terms of the brick-wall cut-off parameter is independent of the horizon surface area. The higher order terms in s are not divergent in h and vanish in the limit $L \rightarrow \infty$. We can extend the above discussions to massive fields. For massive fields $k(r)$ is given by the following expression:

$$k(r) = \frac{1}{V(r)} \sqrt{E^2 - m^2 V(r) - \frac{V(r)}{r^2} l(l+1)} \quad (118)$$

We first find out the maximum value of the term $[\frac{m^2}{V(r)} + \frac{V(r)}{r^2} l(l+1)]$. Thereafter we determine the maximum allowed value of $l(l+1)$ so that the square-root remains real for all values of r and l . The free energy is given by the following expression:

$$\pi\beta F = -\frac{2}{3}\beta \int \frac{dE}{[\exp(\beta E) - 1]} \int \frac{r^2 dr}{V^2(r)} [\{E^2 - m^2 V(r)\}^{\frac{3}{2}} - \{E^2 - m^2 V(r) - s\frac{V(r)}{r^2} N(N+1)\}^{\frac{3}{2}}] \quad (119)$$

Here N is the maximum allowed value of l . We can now follow the steps similar for the massless fields with E^2 replaced by $E^2 - m^2 V(r)$. The leading divergence of the entropy in the brick wall cut-off parameter remains to be of the same form as that for the massless field. The energy is defined with respect to an asymptotic observer and the lower value of the limit of energy integral is m .

To conclude, we have considered the entropy of a scalar field in a Schwarzschild black hole background. We have followed the procedure formulated by 't Hooft. The scalar field is vanishing near the horizon as well as far away from the horizon. The density of states is calculated using the WKB approximation. We found that the entropy associated with the states for which the WKB quantization rule is valid throughout the region of interest is different from that obtained by 't Hooft and as is discussed in the standard literature. The laws of black hole thermodynamics are differential laws. The entropy of the matter fields may be important to interpret the entropy renormalization constant [29]. The part of the entropy associated with the brick-wall cut-off parameter is independent of the black-hole mass and can be considered to give the entropy renormalization constant. However this term is divergent in the brick-wall cut-off parameter and leads to continuous spectrum. Also the universality of the part of the entropy associated with the brick-wall cut-off

parameter indicates the ill-behavedness of this part of the entropy is associated with the intersection of the costant-time foliations at the horizon as is mentioned earlier in this section. In this context it is interesting to study the corresponding situations with other black holes backgrounds. The entropy of the matter fields is also important in the scattering matrix approach of 't Hooft where the entropy of the black hole may be related with the states of the matter fields outside the black hole [29]. This may be significant to explain the black hole entropy if we consider the black hole and the surrounding matter fields outside the horizon as a single system. However we have found in this article that the matter field entropy is not proportional to the horizon surface area, as was obtained in the earlier works, and the above interpretation is not valid for the entropy of the matter fields in the states considered by 't Hooft [29] and in this article. It is also important to note that the entropy of the matter fields in the Euclidian sector is proportional to the area of the horizon and differs from the Lorentzian sector expression obtained in this article. The expression of the matter field entropy for the states considered in this article is also important for the entanglement entropy approach to explain the black hole entropy [44]. The fact that the solutions of the matter field may not contain a global oscillatory term for all values of l' is also important to discuss the scattering and absorption of matter fields in black hole backgrounds. These discussions may also be significant in the context of quasinormal modes. We will consider the above aspects later.

XX. SUPPLEMENT X: RADIATION FROM AN OSCILLATING ELECTRIC DIPOLE

In this section we will consider the solution of the electromagnetic wave equations in the spherical polar coordinates for time varying sources. We will derive an expression for the the electric and magnetic potential of an oscillating point charge and an oscillating electric dipole in the spherical polar coordinates.

In the conventional literature the potentials and the fields in the spherical polar coordinates are assumed to satisfy the non-homogeneous equations similar to the Carteizian coordinates. In the Lorentz gauge the potentials satisfy the non-homogeneous wave equations. The Green's function in the spherical polar coordinates described in reference [23], eqn 6.64, is given by the following expression:

$$G(\vec{R}, \tau) = \frac{1}{2\pi} \int \frac{\exp i[\omega\tau - kR]}{R} d\omega \quad (120)$$

Here $\tau = t - t'$. The ω -integration runs from $-\infty$ to $+\infty$ and k is always positive. However similar to the static situations, in terms of (r, θ, ϕ) the above equation does not satisfy either the homogeneous wave equation or the corresponding Green's function equation. It is easy to find that $\nabla_{\vec{r}}^2 G - \frac{1}{c^2} \frac{\partial^2 G}{\partial t^2}$ contains terms of the form $\int f(R) \frac{\exp i[\omega\tau - kR]}{R} d\omega$ leading to non-vanishing retarded terms in the wave equations. For regular sources the above Green's function leads to the well-known retarded potential:

$$V(\vec{r}, t) = \int \frac{\rho(\vec{r}', t_r) dv'}{R} \quad (121)$$

Where $t_r = t - \frac{R}{c}$ is the retarded time. We can also show from this expression that in terms of (r, θ, ϕ) , the potential does not satisfy the either the homogeneous or the inhomogeneous wave equation thus violating the consistency of the above solution. Only in the Carteizian coordinate the retarded potential satisfies the inhomogeneous wave equation.

We now construct a solution of the Maxwell's equations in the spherical polar coordinates. We follow the discussions made in the context of the static situations. The Maxwell's equations in the spherical polar coordinates defining the fields in terms of the potentials are given by the following homogeneous equations:

$$\vec{\nabla} \cdot \vec{B} = 0 \quad (122)$$

and

$$\vec{\nabla} \times \vec{E} = -\frac{\partial \vec{B}}{\partial t} \quad (123)$$

As we have mentioned in the introduction we here give a potential formulation of electrodynamics in the spherical polar coordinates with time dependent sources. We consider the potentials satisfying the

homogeneous wave equations as fundamental physical variables. This is consistent with a second quantization scheme and as we will observe below give us the experimentally observed static and dynamic potentials and fields. If there exist conjugate pair of harmonics the fields will also satisfy the wave equations and the Lorentz gauge condition will be satisfied by the potentials. However in the spherical polar coordinates this appears not to be the situation. The potentials that we construct from equations (127,128,129..) do not in general satisfy the Lorentz gauge in the spherical polar coordinates and the fields obtained from equation (127,128,129..) may not obey the wave equations in the spherical polar coordinates.

The homogeneous wave equation for the potentials is given by the following expression:

$$\nabla^2 V - \frac{1}{c^2} \frac{\partial^2 V}{\partial t^2} = 0 \quad (124)$$

The most simple solutions of the homogeneous wave equation in the spherical polar coordinates is given by the following expression:

$$g(\vec{r}, t) = N \sum f_l(r, \omega) P_l^m(\cos \gamma) [\exp i\omega t] \quad (125)$$

Here $P_l^m(\cos \gamma)$ are the associated Legendre polynomials and $f_l(r, \omega)$ satisfies the following equation,

$$\frac{\partial}{\partial r} (r^2 \frac{\partial f_l}{\partial r}) - [l(l+1) - \frac{\omega^2}{c^2} r^2] = 0 \quad (126)$$

The construction of the distribution in the spherical polar coordinates is not a straight-forward issue. We have to consider the static distribution to construct the distribution in the dynamic case. Here we consider the $l = 0, 1, 2$ terms for $r > r'$. $f_0(r, \omega)$ is given by the well-known expression:

$$f_0(r, \omega) = a_0 \frac{\exp(-ikr)}{r} \quad (127)$$

$f_1(r, \omega)$ and $f_2(r, \omega)$ are given by the following expressions:

$$f_1(r, \omega) = a_1 [ik + \frac{1}{r}] \frac{\exp(-ikr)}{r} \quad (128)$$

and

$$f_2(r, \omega) = a_2 [-\frac{k^2}{3} + \frac{ik}{r} + \frac{1}{r^2}] \frac{\exp(-ikr)}{r} \quad (129)$$

Here $a_1, a_2, \dots$ are constants. These constants are determined in terms of the sources and the resulting expressions give us the potentials in the dynamical situations. With the normal modes given by the above expressions we can determine the coefficients $a_1, a_2, \dots$ from the static limit of the above solutions. In the static limit the sources are time-independent and $c \rightarrow \infty$. In the static limit the above solutions reduce to expressions similar to the static multipole potentials if we choose the coefficients $a_1, a_2, \dots$ properly as is demonstrated in the following discussions in this section. This is also the situation for points inside source, $r < r'$. Although the wave equation for the potential reduce to the Laplace's equation in the static limit the solutions of the wave equation may not always reduce to the solution of the Laplace's equation in the static limit. We will discuss this situation later and now consider sources for which the above solutions reduce to the corresponding static solutions in the static limit. For localized sources varying periodically with respect to time we define the coefficients $a_1, a_2, \dots$ in terms of the static multipole coefficients determined from the amplitudes of the sinusoidally varying components. To illustrate, in the following we will be considering the radiation from an oscillating electric dipole. In this case we chose $a_0 = q$ and $a_1 = p$ where q is the charges of the oscillating point charges and p is the amplitude of the oscillating electric dipole moment. We also chose $N = \frac{1}{4\pi\epsilon_0}$. The angular frequencies of the radiation are determined from the source frequencies and the corresponding wave numbers are determined from the relation $\omega = ck$ as is valid in the vacuum.

The $\frac{1}{r^2}$ electric field obtained from the above potentials in general does not satisfy the wave equation in the spherical polar coordinates. This is easy to obtain if we consider the angular dependence of the solutions

of the wave equations as are given above. Thus the potentials and the electromagnetic fields are in general not conjugate harmonics in the spherical polar coordinates.

For points inside the source the solutions are given by different expressions and satisfy the homogeneous wave equations with velocity dependent on the permittivity and permeability of the medium. This is similar to the corresponding static situation in the spherical polar coordinates. We will later discuss the solutions of the homogeneous wave equation in the spherical polar coordinates.

We will now consider the potentials of an oscillating point charge and an oscillating electric dipole in the spherical polar coordinates for points outside the source. We have two point charges $q, -q$ oscillating along the Z -axis with the coordinates given by the following expressions:

$$\vec{s}_+ = \frac{s}{2} \cos(\tilde{\omega}t') \hat{k}; \quad \vec{s}_- = -\frac{s}{2} \cos(\tilde{\omega}t') \hat{k} \quad (130)$$

The dipole moment amplitude is $p = qs$. The charge densities are of the form $\rho = q\delta^3(\vec{r}' - \frac{s}{2} \cos(\tilde{\omega}t') \hat{k})$.

In the spherical polar coordinates the potentials of the oscillating point charges that are obtained from the Green's function (120) are given by expressions similar as below:

$$V = \frac{q}{4\pi\epsilon_0} \frac{1}{(1 - \beta^2)^{\frac{1}{2}} |\vec{r}' - \vec{r}'(t'_R)|} \quad (131)$$

Here (t'_R) is the retarded source time: $t'_R = t - \frac{R}{c}$. However the dipole potential obtained from the Green's function (120) in the spherical polar coordinates does not give $(\frac{1}{r})$ radiation fields. On the other hand we will find that the distributions constructed in this article give us $(\frac{1}{r})$ radiation fields for an oscillating electric dipole.

In the Carteizian coordinates the retarded potential due to the oscillating positive charge, obtained from the Lienard-Wiechert potential, is given by the following expression:

$$V(\vec{r}, t) = \frac{q}{4\pi\epsilon_0} \frac{1}{(1 - \vec{\beta} \cdot \hat{R}) |\vec{r} - \vec{s}_+(t_R)|} \quad (132)$$

where $t_R = t - \frac{|\vec{r} - \vec{s}_+(t_R)|}{c}$, $\beta = \frac{v}{c}$. The potential for the negative charge is given by a similar expression. The complete expression in the above equation does not satisfy both the homogeneous and inhomogeneous wave equation in the spherical polar coordinates if we replace the Carteizian coordinates by the spherical polar coordinates. This is similar to the corresponding electrostatic situation. However in the limit of small source velocities, in the dipole approximation $\lambda \gg s$ and for $(x, y, z) \gg (x', y', z')$ the total potentials to the linear order in β and $(\frac{p}{r})$ is similar to the corresponding expressions that are obtained from distributions (127,128,129..) if we replace the Carteizian coordinates by the spherical polar coordinates. The total radiation potential in terms the spherical polar coordinates and linear in the dipole moment is given in equation 9.24 [24]. It may be so that the Lienard-Wiechert potential for an oscillating point charge does not obey the wave equations. We will later discuss the derivation of the Green's function of the wave equation in the Carteizian coordinates given in [23]. It is more appropriate to discuss the radiation problems using the quantum theory of fields.

In the spherical polar coordinates the relationship between the field coordinates and the source coordinates is given by equations (127,128,129..) and also by the higher order solutions. The potentials obey the wave equation by definition.

We will first find the potentials obtained from the monopole term $f_0(r)$. The corresponding potential for an oscillating point charge is given by the following expression:

$$V_0 = \text{Re}[\frac{q}{4\pi\epsilon_0} \frac{1}{r} \exp(\tilde{\omega}t - \tilde{k}r)] \quad (133)$$

This term gives us the static monopole potential in the static limit. In the static limit we recover the Laplace's equation valid for a point charge not at the origin. With an oscillating point charge the right hand side of the wave equation of the monopole potential vanishes for points outside the source $r > s$. For points $r < s$ the distribution will be different. This is similar to the corresponding static potential in the spherical polar coordinates of a point charge not at the origin. The static monopole potential is vanishing for an oscillating electric dipole. We should note that if the point charge is oscillating with a sine function

then the corresponding monopole dynamical potential is vanishing in the static limit and the dynamical solutions do not reduce to the static solution. For a point charge oscillating with the cosine function in eqn (130) replaced by sine function this expected also from the fact that in the static limit the potential of a point charge at the origin satisfy the inhomogeneous Poisson's equation. However in this case also we define the coefficients $a_1, a_2, \dots$ in terms of multipole moments determined from the amplitudes of the sinusoidal oscillations.

We now consider $(\frac{1}{r})$ radiation potentials. These terms are obtained from the $(\frac{1}{r})$ terms present in the normal modes $f_l(r, \tilde{k})$ with $l \geq 1$. The $(\frac{1}{r})$ radiation potential linear in the dipole moment is obtained from f_1 and is given by the following expression:

$$V = \text{Re}\left\{\frac{i}{4\pi\epsilon_0} \frac{p\tilde{\omega}}{cr} \exp[\tilde{\omega}t - \tilde{k}r](\cos\theta)\right\} \quad (134)$$

With the assumed form of the time varying sources the electric potential is given by the following expression:

$$V = -\frac{1}{4\pi\epsilon_0}(\cos\theta) \frac{p\tilde{\omega}}{c} \frac{\sin(\tilde{\omega}t - \tilde{k}r)}{r} \quad (135)$$

Here $\vec{k} = \tilde{k}\hat{r}$ and $\tilde{\omega} = c\tilde{k}$. The above expression is similar to the corresponding expression obtained in [24]. The second term in f_1 gives us the static dipole electric potential in the static limit of $\tilde{\omega} \rightarrow 0$ and $c \rightarrow \infty$.

The $(\frac{1}{r})$ terms present in the higher order $f_l(r, \tilde{k})$ s also give $(\frac{1}{r})$ radiation fields. However in the perfect dipole radiation limit, $\lambda \gg s$, the amplitudes of these terms are of higher order smallness and we neglect these contributions.

The current density is given by $\vec{J} = \rho\vec{v}$. Each component of the magnetic vector potential satisfies the homogeneous wave equation in the spherical polar coordinates. We can obtain the magnetic vector potential following the above procedure with the coefficients $a_1, a_2, \dots$ determined in terms of the magnetostatic multipole coefficients. To the leading order in the dipole moment p the radiation vector potential is obtained from $f_0(r)$ term and is given by the following expression:

$$\vec{A} = -\frac{\mu_0}{4\pi} p\tilde{\omega} \frac{\sin(\tilde{\omega}t - \tilde{k}r)}{r} \hat{z} \quad (136)$$

The higher order terms in the distributions (127,128,129..) do not contribute radiation terms linear in p in the oscillating electric dipole magnetic potential. The above expression is similar to the corresponding expression obtained in [24].

We now consider the derivation of the electric and magnetic fields from the potentials. We take the derivatives of the potentials using the Cartesian coordinate unit vectors and transform the unit vectors in the resulting fields to the spherical polar unit vectors. The resulting electric and the magnetic fields, to the order linear in $(\frac{p\omega^2}{r})$, are given by the following expressions:

$$\vec{E} = -\frac{\mu_0 p \omega^2}{4\pi r} (\sin\theta) \sin(\omega t - kr) \hat{\theta} \quad (137)$$

and,

$$\vec{B} = -\frac{\mu_0 p \omega^2}{4\pi cr} (\sin\theta) \sin(\omega t - kr) \hat{\phi} \quad (138)$$

Here ω is the angular frequency of the harmonic oscillator. The electromagnetic fields give the forces acting on charged particles and currents. However the electromagnetic fields in the spherical polar coordinates obtained from equations (127..) do not obey the wave equations and are not conjugate harmonics of the corresponding potentials. We will later discuss the polarization properties of the electromagnetic fields.

We now consider the power emitted by the oscillating electric dipole. The Poynting's vector is to be obtained from the free field Lagrangian expressed in terms of the four potentials $A_\mu = (\frac{V}{c}, \vec{A})$. The Lagrangian density is given by the expression: $-\frac{c}{\mu_0} A_{\alpha,\beta} A_{\alpha,\beta}$. The time averaged radial power obtained from the potentials (135) and (136) is given by the following expression [47]:

$$\langle \vec{S} \rangle = \frac{\mu_0}{2c} \left[\frac{p\omega^2}{4\pi r} \right]^2 [(\sin \theta)^2] \hat{r} \quad (139)$$

Thus the time-averaged radial power, the expression analogous to the radial Poynting vector, coincides with the conventional expression [24]:

$$\langle \vec{S}' \rangle = \frac{\mu_0}{2c} \left[\frac{p\omega^2}{4\pi r} \right]^2 [(\sin \theta)^2] \hat{r} \quad (140)$$

However the time-dependent energy-momentum tensor may be different from the conventional expressions [24]. This is expected from the fact that in general, in an arbitrary curvilinear coordinate system, the potentials may not satisfy both the wave equations and the Lorentz gauge condition. It will be convenient to consider the potentials from the individual oscillating charges to verify that the potentials give a conserved energy-momentum four vector. We will later discuss this issue. Equation (140) is obtained by replacing the Cartesian coordinates (x, y, z) by the spherical polar coordinates (r, θ, ϕ) in the expressions for the potentials obtained from the Lienard-Weichert potential. We have mentioned earlier that the potentials so obtained do not satisfy the wave equations in the spherical polar coordinates.

An interesting aspect of the potential of an oscillating point charge obtained from the distributions (127...) is the following. The $(\frac{1}{r})$ terms in the higher order normal modes $f_l(r, r')$ contributes $(\frac{1}{r})$ radiation terms to the electric potential with angular frequency ω where ω is the frequency of the oscillating point charge. Similar discussions are also valid for the corresponding magnetic vector potential. In the dipole approximation the amplitudes of these terms are of higher order smallness in the parameter (ks) .

We now construct the solution of the homogeneous wave equation with $r < r'$. Where (r) is the radial coordinate of the field point and (r') is the radial coordinate of the source point. Considering the solutions of the static Laplace's equation and the solutions of the wave equation for $r > r'$ we assume that the radial solutions of the wave equation for $r < r'$ are given by expressions of the following form:

$$\chi_l = \alpha(r, l) r^l \exp(ikr) \quad (141)$$

If we substitute the above expression in the radial part of the wave equation we have the following equation for the function $\alpha(r, l)$:

$$r\alpha'' + [2(l+1) + 2ikr]\alpha' + 2ik(l+1)\alpha = 0 \quad (142)$$

It is easy to solve the above equation and the complete potential for $r < r'$ is given by the real or imaginary part of the following expression depending on the nature of the source:

$$R(\vec{r}, t) = \sum_l \left[\sum_j a_j \frac{r^{(l+j)}}{r'^{(l+1)}} P_l(\cos \gamma) \right] \exp[i(\omega t + kr)] \quad (143)$$

Here the summations over (l) and (j) are running from zero to infinity and if we solve the indicial equation [46] for arbitrary values l together with the required regularity conditions at the origin we have the following recurrence relation,

$$a_{j+1} = -(2ik) \left[\frac{(l+j+1)}{(j+1)(j+2l+2)} \right] a_j \quad (144)$$

we chose N to be given by the following expression:

$$N = \frac{1}{4\pi\epsilon_0} \quad (145)$$

The constants $a_0(l)$ are determined from the source similar to the situation of $r > r'$. For a source oscillating according to a cosine function of time the above solution reduce to the corresponding static solution

of the Laplace's equation in the static limit of static source and $c \rightarrow \infty$ if we choose $a_0(l) = \int \rho_0(\vec{r}') dv'$ and put the $\vec{r}'$ dependent terms into the integral. Here $\rho_0(\vec{r}')$ is the amplitude of the oscillating source.

We now demonstrate that in terms of an arbitrary curvilinear coordinate system it is in general not possible to find a gauge so that the potentials satisfy both the wave equations and the Lorentz gauge condition. Let A^μ be a set of potentials which satisfy the wave equation but $\partial_\mu A^\mu \neq 0$. If there exists a function λ such that $A^\mu + \partial^\mu \lambda$ satisfy both the Lorentz gauge condition and the wave equations then we have the following two conditions on λ respectively:

$$\nabla^2 \lambda - \frac{1}{c^2} \frac{\partial^2 \lambda}{\partial t^2} = -\partial_\mu A^\mu \quad (146)$$

and

$$\partial_t [\nabla^2 \lambda - \frac{1}{c^2} \frac{\partial^2 \lambda}{\partial t^2}] = 0 \quad (147)$$

The above two conditions are obtained from the Lorentz gauge condition and from the fact that the electric potential satisfies the wave equations. The above two conditions are in general contradictory. This happens to be the situation with the fields of an oscillating electric dipole.

In this section we have only considered the radiation field from oscillating charges. A more complete description of radiation problems can be possible through an appropriate quantum theory with the distributions determined from a quantum theory. The discussions in this article may also indicate that in all physically allowed space-time there exist a preferred coordinate system where the Poisson's equation is satisfied for a point charge and also the electromagnetic potentials and the electromagnetic fields are conjugate harmonics to each other in four dimension in the Lorentz gauge. This appears to be the situation in the flat space-time with the Cartesian coordinate system.

XXI. MISCELLANEOUS

One can observe Newton's rings with plane-surface of the convex lens on the glass plate.

We now discuss an elementary method to calculate the surface integral of a vector field. Let us consider the plane $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$ in the positive octant. We consider curves on the plane parallel to the $X - Z$ plane. The curves are given by the following equation:

$$\frac{x}{a} + \frac{z}{c} = 1 - \frac{y}{b} \quad (148)$$

where y acts as a parameter. An infinitesimal line element on these curves are given by $d\vec{u} = (dx)\hat{i} - (\frac{c}{a})(dx)\hat{k}$. Similarly we can consider curves on the plane parallel to the $Y - Z$ plane. The curves are given by the following equation:

$$\frac{y}{b} + \frac{z}{c} = 1 - \frac{x}{a} \quad (149)$$

where x acts as a parameter. An infinitesimal line element on these curves are given by $d\vec{v} = (dy)\hat{j} - (\frac{c}{b})(dy)\hat{k}$. An area element on the plane is given by $d\vec{u} \times d\vec{v}$ and is expressed in terms of $dx dy$. We can thereafter do the surface integral. One can generalize this method properly. These discussions may also be relevant to find out whether a scalar field has an equipotential surface.

In the following section we will describe the force configuration acting on a uniform cube with weight ' W ' resting on a horizontal plane with friction. The coefficient of static friction is ' μ ' and the sides of the cube are of length ' $2a$ '. We consider a coordinate system on the surface with the origin, O , directly below the center of the cube. Let a force ' $\vec{P}$ ' act on the block which is not large enough to move the block. The force acts along the X -axis. Since the block is not moving the friction between the surfaces should develop an equal and opposite force ' $-\vec{P}$ ' acting on the cube at the interface balancing ' $\vec{P}$ '. However if we consider the torque balance w.r.t the origin then the reaction force at the interface will not be uniform. The reaction force per unit area ' $\vec{R}$ ' becomes a function of x such that:

$$Pa = 2a \int_{-a}^a xR(x)dx \quad (150)$$

The weight is given by a surface integral of $R(x)$ and we have the following two equations:

$$P = 2 \int_{-a}^a xR(x)dx \quad (151)$$

and

$$W = 2a \int_{-a}^a R(x)dx \quad (152)$$

It is expected that $R(x)$ will be of the form: $R + r(x, P)$, where R is constant so that the surface integral of R is W and $r(x, P)$ is an antisymmetric function w.r.t x such that the equation just above remains valid. When P is large enough so that the block is on the verge of motion we have the following relation:

$$P_{th} = 2\mu a \int_{-a}^a R_{th}(x)dx \quad (153)$$

The first equation then leads to the following condition on $r(x, P)$:

$$\mu W = 2 \int_{-a}^a xr(x, P_{th})dx \quad (154)$$

which may be useful to determine $r(x, P_{th})$ if we consider linear relationships.

XXII. END

Project Euclidean (Section:II) is complete.

Project Galileo-Maxwell (Supplement:II,III) is complete.

XXIII. ACKNOWLEDGMENT AND

I am thankful to Prof. D. J. Griffiths for pointing out a correction regarding equ.(51). I am thankful to a member of VECC Housing, Kolkata for reminding me an important point. I am also thankful to a few faculty members and of B-12 (2006-07) students of HIT.

XXIV. REFERENCES

- [1] J. J. Sakurai; Advanced Quantum Mechanics: Addison-Wesley Publishing Company.
- [2] A. N. Matveev; Mechanics and Theory of Relativity: Mir Publishers, Moscow.
- [3] R. M. Wald; General Relativity: The University of Chicago Press, Chicago and London.
- [4] C. W. Misner, K. S. Thorne, J. A. Wheeler; Gravitation: W. H. Freeman and Company, New York.
- [5] A. N. Kolmogorov and S. V. Fomin; Functional Analysis (Vol. I): Dover Publications, Inc. New York.
- [6] W. Rudin; Introduction to Topology and Modern Analysis: Mc Graw - Hill Book Company.
- [7] W. Lamb (Jr.); Physics Today 22.
- [8] S. Chandrasekhar.
- [9] A. K. Raychoudhury.
- [10] N. N. Bogoliubov, D. V. Shirkov; Introduction To The Theory of Quantized Fields: John Wiley and sons.
- [11] A. K. Raychoudhury *et al* .

- [12] A. Afriat, F. Sellery; The Einstein, Podolsky and Rosen paradox: Plenum publishing corporation.
- [13] J. J. Sakurai; Modern Quantum Mechanics (Section: 1.4 - 1.6): Addison Wesley Publishing Company.
- [14] K. Ghosh; gr-qc/0504083, gr-qc/0503003.
- [15] M. Nakahara; Geometry, Topology And Physics: Adam Hilger, Bristol And New York.
- [16] T. M. Apostol; Mathematical Analysis: Narosa Publishing House.
- [17] S. Weinberg; Gravitation and Cosmology: John Wiley and Sons.
- [18] M. E. Preskin, D. V. Schroeder; An Introduction to Quantum Field Theory: Addison-Wesley Publishing Company.
- [19] S. W. Hawking, G. F. R. Ellis; The Large Scale Structure of Space-Time: Cambridge University Press.
- [20] L. D. Landau and E. M. Lifshitz; The Classical Theory of Fields: Butterworth-Heinemann.
- [21] J. D. Bjorken, S. D. Drell; Relativistic Quantum Fields: Mc Graw-Hill Book Company.
- [22] P. Ramond; Field Theory: A Modern Primer, Addison Wesley Publishing Company
- [23] J. D. Jackson; Classical Electrodynamics: Wiley Eastern Limited.
- [24] D. J. Griffiths; Introduction To Electrodynamics(1989): Prentice-Hall of India.
- [25] R. P. Feynman, R. B. Leighton, M. Sands: The Feynman Lectures on Physics; Narosa Publishing House.
- [26] F. A. Jenkins, H. E. Harvey; Fundamentals of Optics: McGraw-Hill Book Company.
- [27] R. Longhurst; Geometric and Physical Optics: Orient-Longman.
- [28] G. B. Arfken, H. J. Weber; Mathematical Methods For Physicists: Academic Press, Inc.
- [29] G. 't Hooft, Nucl. Phys. B 256, (1985)
- [30] F. K. Richtmyer, E. H. Kennard and J. N. Cooper; Introduction to Modern Physics: Tata McGraw-Hill Publishing Company, New Delhi.
- [31] A. Beiser; Concepts of Modern physics: McGraw-Hill International Editions (1987).
- [32] W. K. Clifford: The Postulates of the Science of Space, The World of Mathematics: Vol.1; Editor: James Newman, Dover Publications, Inc.
- [33] R. Courant and F. John, Introduction to Calculus and Analysis: Vol.1; Springer.
- [34] L. Carroll: Continuity, The World of Mathematics: Vol.4; Editor: James Newman, Dover Publications, Inc.
- [35] Wikipedia, The Free Encyclopedia.
- [36] R. Dedekind: Irrational Numbers, The World of Mathematics: Vol.1; Editor: James Newman, Dover Publications, Inc.
- [37] R. Resnick; Introduction to Special Relativity: Wiley Eastern Limited.
- [38] C. Itzykson and J. B. Zuber; Quantum Field Theory: McGraw-Hill Book Company (1986).
- [39] J. Bekenstein, Phys. Rev. D 7, (1973)
- [40] S. W. Hawking, Commun. Math. Phys. 43, (1975)
- [41] L. Susskind and J. Uglum, Phys. Rev. D 50, 2700, (1994)
- [42] A. Ghosh and P. Mitra, Phys. Lett. B 357, 295, (1995)
- [43] I. Ichinose and Y. Satoh, Nucl. Phys. B 447, 34, (1995)
- [44] L. Bombelli, R. K. Kaul, J. Lee and R. Sorkin, Phys. Rev. D 34, 373 (1986)
- [45] A. Beiser, Concepts of Modern Physics: McGraw-Hill Book Company.
- [46] M. R. Spiegel and A. N. Pettitt, Advanced Mathematics for Engineers and Scientists: McGraw-Hill International Book Company.
- [47] H. Goldstein, Classical Mechanics: Narosa Publishing House.